This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



A2	-
	(43) International Publication Date: 18 May 2000 (18.05.0
	BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, G
: THI SYLVA	ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SS, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KMD, RU, TJ, TM), European patent (AT, BE, CH, CY, EDK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SIOAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, MC, CH, CH, CH, CH, CH, CH, CH, CH, CH, C
A 1903:	upon receipt of that report.
Howson, Spring	
I NUCL	EIC ACID SEQUENCES, VECTORS AND HOST CELLS CONTAI
(AAV) are met	serotype 1 are provided, as are vectors and host cells containing the nods of delivering genes via AAV-1 derived vectors.
) : 5 6	mes, M. A 19035 S Wash-Howson, Spring

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	Fl	Finland	LT	Lithuania	SK	Slovakia
АT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	T.J	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
СН	Switzerland	KG	Kyrgyzstan -	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand	2011	Zillioabwc
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		•
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	Li	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

ADENO-ASSOCIATED VIRUS SEROTYPE I NUCLEIC ACID SEQUENCES, VECTORS AND HOST CELLS CONTAINING SAME

This work was supported by the National Institutes of Health, grant no. P30 DK47757-06 and PO1 HD32649-04. The US government may have certain rights in this invention.

Field of the Invention

5

10

15

20

25

This invention relates generally to viral vector, and more particularly, to recombinant viral vectors useful for gene delivery.

Background of the Invention

Adeno-associated viruses are small, single-stranded DNA viruses which require helper virus to facilitate efficient replication [K.I. Berns, Parvoviridae: the viruses and their replication, p. 1007-1041, in F.N. Fields et al., Fundamental virology, 3rd ed., vol. 2, (Lippencott-Raven Publishers, Philadelphia, PA) (1995)]. The 4.7 kb genome of AAV is characterized by two inverted terminal repeats (ITR) and two open reading frames which encode the Rep proteins and Cap proteins, respectively. The Rep reading frame encodes four proteins of molecular weight 78 kD, 68 kD, 52 kD and 40 kD. These proteins function mainly in regulating AAV replication and integration of the AAV into a host cell's chromosomes. The Cap reading frame encodes three structural proteins in molecular weight 85 kD (VP 1), 72 kD (VP2) and 61 kD (VP3) [Berns, cited above]. More than 80% of total proteins in AAV virion comprise VP3. The two ITRs are the only cis elements essential for AAV replication, packaging and integration. There are two conformations of AAV ITRs called "flip" and "flop". These differences in conformation originated from the replication model of adeno-associated virus which use the ITR to initiate and reinitiate the replication [R.O. Snyder et al., J. Virol., 67:6096-6104 (1993), K.I. Berns, Microbiological Reviews, 54:316-329 (1990)].

AAVs have been found in many animal species, including primates, canine, fowl and human [F.A. Murphy et al., "The Classification and Nomenclature of Viruses: Sixth Report of the International Committee on Taxonomy of Viruses".

10

Archives of Virology, (Springer-Verlag, Vienna) (1995)]. In addition to five known primate AAVs (AAV-1 to AAV-5), AAV-6, another serotype closely related to AAV-2 and AAV-1 has also been isolated [E. A. Rutledge et al., J. Virol., 72:309-319 (1998)]. Among all known AAV serotypes, AAV-2 is perhaps the most well-characterized serotype, because its infectious clone was the first made [R.J. Samulski et al., Proc. Natl. Acad. Sci. USA, 79:2077-2081 (1982)]. Subsequently, the full sequences for AAV-3A, AAV-3B, AAV-4 and AAV-6 have also been determined [Rutledge, cited above; J.A.Chiorini et al., J. Virol., 71:6823-6833 (1997); S. Muramatsu et al., Virol., 221:208-217 (1996)]. Generally, all AAVs share more than 80% homology in nucleotide sequence.

A number of unique properties make AAV a promising vector for human gene therapy [Muzyczka, Current Topics in Microbiology and Immunology, 158:97-129 (1992)]. Unlike other viral vectors, AAVs have not been shown to be associated with any known human disease and are generally not considered pathogenic. Wild type 15 AAV is capable of integrating into host chromosomes in a site specific manner [R. M. Kotin et al., Proc. Natl. Acad. Sci. USA, 87:2211-2215 (1990)- R.J. Samulski, EMBO J., 10(12):3941-3950 (1991)]. Recombinant AAV vectors can integrate into tissue cultured cells in chromosome 19 if the rep proteins are supplied in trans [C. Balague et al., J. Virol., 71:3299-3306 (1997); R. T. Surosky et al., J. Virol., 71:7951-7959 (1997)]. The integrated genomes of AAV have been shown to allow 20 long term gene expression in a number of tissues, including, muscle, liver, and brain [K. J. Fisher, Nature Med., 3(3):306-312 (1997); R. O. Snyder et al., Nature Genetics, 16:270-276 (1997); X. Xiao et al., Experimental Neurology, 144:113-124 (1997); Xiao, J. Virol., 70(11):8098-8108 (1996)].

AAV-2 has been shown to be present in about 80-90% of the human population. Earlier studies showed that neutralizing antibodies for AAV-2 are prevalent [W. P. Parks et al., <u>J. Virol.</u>, <u>2</u>:716-722 (1970)]. The presence of such antibodies may significantly decrease the usefulness of AAV vectors based on AAV-2 despite its other merits. What are needed in the art are vectors characterized by the

10

15

20

25

advantages of AAV-2, including those described above, without the disadvantages, including the presence of neutralizing antibodies.

Summary of the Invention

In one aspect, the invention provides an isolated AAV-1 nucleic acid molecule which is selected from among SEQ ID NO: 1, the strand complementary to SEQ ID NO: 1, and cDNA and RNA sequences complementary to SEQ ID NO: 1 and its complementary strand.

In another aspect, the present invention provides AAV ITR sequences, which include the 5' ITR sequences, nt 1 to 143 of SEQ ID NO: 1; the 3' ITR sequences, nt 4576 to 4718 of SEQ ID NO: 1, and fragments thereof.

In yet another aspect, the present invention provides a recombinant vector comprising an AAV-1 ITR and a selected transgene. Preferably, the vector comprises both the 5' and 3' AAV-1 ITRs between which the selected transgene is located.

In still another aspect, the invention provides a recombinant vector comprising an AAV-1 P5 promoter having the sequence of nt 236 to 299 of SEQ ID NO: 1 or a functional fragment thereof.

In a further aspect, the present invention provides a nucleic acid molecule encoding an AAV-1 rep coding region and an AAV-1 cap coding region.

In still another aspect, the present invention provides a host cell transduced with a recombinant viral vector of the invention. The invention further provides a host cell stably transduced with an AAV-1 P5 promoter of the invention.

In still a further aspect, the present invention provides a pharmaceutical composition comprising a carrier and a vector of the invention.

In yet another aspect, the present invention provides a method for AAV-mediated delivery of a transgene to a host involving the step of delivering to a selected
host a recombinant viral vector comprising a selected transgene under the control of
sequences which direct expression thereof and an adeno-associated virus 1 (AAV-1)
virion.

15

20

25

In another aspect, the invention provides a method for in vitro production of a selected gene product using a vector of the invention.

Other aspects and advantages of the invention will be readily apparent to one of skill in the art from the detailed description of the invention.

5 Brief Description of the Drawings

Figs. 1A-1C illustrate the alignment of nucleotides of AAV-1 [SEQ ID NO: 1], AAV-2 [SEQ ID NO: 18] and AAV-6 [SEQ ID NO: 19]. The alignment was done with MacVector 6.0. The full sequences of AAV-1 are shown in the top line. Nucleotides in AAV-2 and AAV-6 identical to AAV-1 are symbolized by "." and gaps by "-". Some of the conserved features among AAVs are marked in this figure. Note the 3' ITRs of AAV-1 and AAV-6 are shown in different orientations.

Fig. 2 illustrates the predicted secondary structure of AAV-1 ITR. The nucleotides in AAV-2 and AAV-6 are shown in italic and bold respectively.

Fig. 3A illustrates a hypothesis of how AAV-6 arose from the homologous recombination between AAV-1 and AAV-2. The major elements of AAV-1 are indicated in the graph. A region that is shared between AAV-1, AAV-2 and AAV-6 is shown in box with waved lines.

Fig. 3B is a detailed illustration of a 71 bp homologous region among AAV-1, AAV-2 and AAV-6. Nucleotides that differ among these serotypes are indicated by arrows.

Fig. 4A is a bar chart illustrating expression levels of human alpha 1 antitrypsin (α1AT) in serum following delivery of hAAT via recombinant AAV-1 and recombinant AAV-2 viruses.

Fig. 4B is a bar chart illustrating expression levels of erythropoietin (epo) in serum following delivery of the epo gene via recombinant AAV-1 and recombinant AAV-2 viruses.

Fig. 5A is a bar chart illustrating expression levels of $\alpha 1AT$ in liver following delivery of $\alpha 1AT$ as described in Example 7.

10

20

25

Fig. 5B is a bar chart demonstrating expression levels of epo in liver following delivery of epo as described in Example 7.

Fig. 5C is a bar chart demonstrating neutralizing antibodies (NAB) directed to AAV-1 following delivery of a1AT or epo to liver as described in Example 7.

Fig. 5D is a bar chart demonstrating neutralizing antibodies (NAB) directed to AAV-2 following delivery of α1AT or epo to liver as described in Example 7.

Fig. 6A is a bar chart illustrating expression levels of α1AT in muscle following delivery of α1AT as described in Example 7.

Fig. 6B is a bar chart demonstrating expression levels of epo in muscle following delivery of epo as described in Example 7.

Fig. 6C is a bar chart demonstrating neutralizing antibodies (NAB) directed to AAV-1 following delivery of a1AT or epo to muscle as described in Example 7.

Fig. 6D is a bar chart demonstrating neutralizing antibodies (NAB) directed to AAV-2 following delivery of a1AT or epo to muscle as described in Example 7.

15 Detailed Description of the Invention

The present invention provides novel nucleic acid sequences for an adeno-associated virus of serotype 1 (AAV-1). Also provided are fragments of these AAV-1 sequences. Among particularly desirable AAV-1 fragments are the inverted terminal repeat sequences (ITRs), rep and cap. Each of these fragments may be readily utilized, e.g., as a cassette, in a variety of vector systems and host cells. Such fragments may be used alone, in combination with other AAV-1 sequences or fragments, or in combination with elements from other AAV or non-AAV viral sequences. In one particularly desirable embodiment, a cassette may contain the AAV-1 ITRs of the invention flanking a selected transgene. In another desirable embodiment, a cassette may contain the AAV-1 rep and/or cap proteins, e.g., for use in producing recombinant (rAAV) virus.

Thus, the AAV-1 sequences and fragments thereof are useful in production of rAAV, and are also useful as antisense delivery vectors, gene therapy vectors, or vaccine vectors. The invention further provides nucleic acid molecules, gene delivery

10

15

20

25

vectors, and host cells which contain the AAV-1 sequences of the invention. Also provided a novel methods of gene delivery using AAV vectors.

As described herein, the vectors of the invention containing the AAV-1 capsid proteins of the invention are particularly well suited for use in applications in which the neutralizing antibodies diminish the effectiveness of other AAV serotype based vectors, as well as other viral vectors. The rAAV vectors of the invention are particularly advantageous in rAAV readministration and repeat gene therapy.

These and other embodiments and advantages of the invention are described in more detail below. As used throughout this specification and the claims, the term "comprising" is inclusive of other components, elements, integers, steps and the like.

I. AAV-1 NUCLEIC ACID AND PROTEIN SEQUENCES

The AAV-1 nucleic acid sequences of the invention include the DNA sequences of SEQ ID NO: 1 (Figs. 1A-1C), which consists of 4718 nucleotides. The AAV-1 nucleic acid sequences of the invention further encompass the strand which is complementary to SEQ ID NO: 1, as well as the RNA and cDNA sequences corresponding to SEQ ID NO: 1 and its complementary strand. Also included in the nucleic acid sequences of the invention are natural variants and engineered modifications of SEQ ID NO: 1 and its complementary strand. Such modifications include, for example, labels which are known in the art, methylation, and substitution of one or more of the naturally occurring nucleotides with an analog.

Further included in this invention are nucleic acid sequences which are greater than 85%, preferably at least about 90%, more preferably at least about 95%, and most preferably at least about 98 - 99% identical or homologous to SEQ ID NO.1. The term "percent sequence identity" or "identical" in the context of nucleic acid sequences refers to the residues in the two sequences which are the same when aligned for maximum correspondence. The length of sequence identity comparison may be over the full-length sequence, or a fragment at least about nine nucleotides, usually at least about 20 - 24 nucleotides, at least about 28 - 32 nucleotides, and preferably at least about 36 or more nucleotides. There are a number of different

WO 00/28061

5

10

7

algorithms known in the art which can be used to measure nucleotide sequence identity. For instance, polynucleotide sequences can be compared using Fasta, a program in GCG Version 6.1. Fasta provides alignments and percent sequence identity of the regions of the best overlap between the query and search sequences (Pearson, 1990, herein incorporated by reference). For instance, percent sequence identity between nucleic acid sequences can be determined using Fasta with its default parameters (a word size of 6 and the NOPAM factor for the scoring matrix) as provided in GCG Version 6.1, herein incorporated by reference.

The term "substantial homology" or "substantial similarity," when referring to a nucleic acid or fragment thereof, indicates that, when optimally aligned with appropriate nucleotide insertions or deletions with another nucleic acid (or its complementary strand), there is nucleotide sequence identity in at least about 95 - 99% of the sequence.

Also included within the invention are fragments of SEQ ID NO: 1, its 15 complementary strand, cDNA and RNA complementary thereto. Suitable fragments are at least 15 nucleotides in length, and encompass functional fragments which are of biological interest. Certain of these fragments may be identified by reference to Figs. 1A-1C. Examples of particularly desirable functional fragments include the AAV-1 inverted terminal repeat (ITR) sequences of the invention. In contrast to the 145 nt ITRs of AAV-2, AAV-3, and AAV-4, the AAV-1 ITRs have been found to consist of 20 only 143 nucleotides, yet advantageously are characterized by the T-shaped hairpin structure which is believed to be responsible for the ability of the AAV-2 ITRs to direct site-specific integration. In addition, AAV-1 is unique among other AAV serotypes, in that the 5' and 3' ITRs are identical. The full-length 5' ITR sequences of AAV-1 are provided at nucleotides 1-143 of SEQ ID NO: 1 (Fig. 1A) and the full-25 length 3' ITR sequences of AAV-1 are provided at nt 4576-4718 of SEQ ID NO: 1 (Fig. 1C). One of skill in the art can readily utilize less than the full-length 5' and/or 3' ITR sequences for various purposes and may construct modified ITRs using conventional techniques, e.g., as described for AAV-2 ITRs in Samulski et al, Cell, 30 33:135-143 (1983).

10

15

20

25

Another desirable functional fragment of the AAV-1 genome is the P5 promoter of AAV-1 which has sequences unique among AAV P5 promoters, while maintaining critical regulatory elements and functions. This promoter is located within nt 236 - 299 of SEQ ID NO: 1 (Fig. 1A). Other examples of functional fragments of interest include the sequences at the junction of the rep/cap, e.g., the sequences spanning nt 2306-2223, as well as larger fragments which encompass this junction which may comprise 50 nucleotides on either side of this junction. Still other examples of functional fragments include the sequences encoding the rep proteins. Rep 78 is located in the region of nt 334 - 2306 of SEQ ID NO: 1; Rep 68 is located in the region of nt 334-2272, and contains an intron spanning nt 1924-2220 of SEQ ID NO: 1. Rep 52 is located in the region of nt 1007 - 2304 of SEQ ID NO: 1; rep 40 is located in the region of nt 1007 - 2272, and contains an intron spanning nt 1924-2246 of SEQ ID NO: 1. Also of interest are the sequences encoding the capsid proteins, VP 1 [nt 2223-4431 of SEQ ID NO: 1], VP2 [nt 2634-4432 of SEQ ID NO: 1] and VP3 [nt 2829-4432 of SEQ ID NO: 1]. Other fragments of interest may include the AAV-1 P19 sequences, AAV-1 P40 sequences, the rep binding site, and the terminal resolute site (TRS).

The invention further provides the proteins and fragments thereof which are encoded by the AAV-1 nucleic acids of the invention. Particularly desirable proteins include the rep and cap proteins, which are encoded by the nucleotide sequences identified above. These proteins include rep 78 [SEQ ID NO:5], rep 68 [SEQ ID NO:7], rep 52 [SEQ ID NO:9], rep 40 [SEQ ID NO: 11], vpl [SEQ ID NO: 13], vp2 [SEQ ID NO: 15], and vp3 [SEQ IID NO: 17] and functional fragments thereof while the sequences of the rep and cap proteins have been found to be closely related to those of AAV-6, there are differences in the amino acid sequences (see Table 1 below), as well as differences in the recognition of these proteins by the immune system. However, one of skill in the art may readily select other suitable proteins or protein fragments of biological interest. Suitably, such fragments are at least 8 amino acids in length. However, fragments of other desired lengths may be readily utilized.

10

15

20

25

Such fragments may be produced recombinantly or by other suitable means, e.g., chemical synthesis.

The sequences, proteins, and fragments of the invention may be produced by any suitable means, including recombinant production, chemical synthesis, or other synthetic means. Such production methods are within the knowledge of those of skill in the art and are not a limitation of the present invention.

II. VIRAL VECTORS

In another aspect, the present invention provides vectors which utilize the AAV-1 sequences of the invention, including fragments thereof, for delivery of a heterologous gene or other nucleic acid sequences to a target cell. Suitably, these heterologous sequences (i.e., a transgene) encode a protein or gene product which is capable of being expressed in the target cell. Such a transgene may be constructed in the form of a "minigene". Such a "minigene" includes selected heterologous gene sequences and the other regulatory elements necessary to transcribe the gene and express the gene product in a host cell. Thus, the gene sequences are operatively linked to regulatory components in a manner which permit their transcription. Such components include conventional regulatory elements necessary to drive expression of the transgene in a cell containing the viral vector. The minigene may also contain a selected promoter which is linked to the transgene and located, with other regulatory elements, within the selected viral sequences of the recombinant vector.

Selection of the promoter is a routine matter and is not a limitation of this invention. Useful promoters may be constitutive promoters or regulated (inducible) promoters, which will enable control of the timing and amount of the transgene to be expressed. For example, desirable promoters include the cytomegalovirus (CMV) immediate early promoter/enhancer [see, e.g., Boshart et al, Cell, 41:521-530 (1985)], the Rous sarcoma virus LTR promoter/enhancer, and the chicken cytoplasmic β-actin promoter [T. A. Kost et al, Nucl, Acids Res., 11(23):8287 (1983)]. Still other desirable promoters are the albumin promoter and an AAV P5 promoter. Optionally, the selected promoter is used in conjunction with a heterologous enhancer, e.g., the β-

10

actin promoter may be used in conjunction with the CMV enhancer. Yet other suitable or desirable promoters and enhancers may be selected by one of skill in the art.

5

10

15

20

25

30

The minigene may also desirably contain nucleic acid sequences heterologous to the viral vector sequences including sequences providing signals required for efficient polyadenylation of the transcript (poly-A or pA) and introns with functional splice donor and acceptor sites. A common poly-A sequence which is employed in the exemplary vectors of this invention is that derived from the papovavirus SV-40. The poly-A sequence generally is inserted in the minigene downstream of the transgene sequences and upstream of the viral vector sequences. A common intron sequence is also derived from SV-40, and is referred to as the SV40 T intron sequence. A minigene of the present invention may also contain such an intron, desirably located between the promoter/enhancer sequence and the transgene. Selection of these and other common vector elements are conventional [see, e.g., Sambrook et al, "Molecular Cloning. A Laboratory Manual", 2d edit., Cold Spring Harbor Laboratory, New York (1989) and references cited therein] and many such sequences are available from commercial and industrial sources as well as from Genebank.

The selection of the transgene is not a limitation of the present invention. Suitable transgenes may be readily selected from among desirable reporter genes, therapeutic genes, and optionally, genes encoding immunogenic polypeptides. Examples of suitable reporter genes include β-galactosidase (β-gal), an alkaline phosphatase gene, and green fluorescent protein (GFP). Examples of therapeutic genes include, cytokines, growth factors, hormones, and differentiation factors, among others. The transgene may be readily selected by one of skill in the art. See, e.g., WO 98/09657, which identifies other suitable transgenes.

Suitably, the vectors of the invention contain, at a minimum, cassettes which consist of fragments of the AAV-1 sequences and proteins. In one embodiment, a vector of the invention comprises a selected transgene, which is flanked by a 5' ITR and a 3' ITR, at least one of which is an AAV-1 ITR of the invention. Suitably,

11

vectors of the invention may contain a AAV-1 P5 promoter of the invention. In yet another embodiment, a plasmid or vector of the invention contains AAV-1 rep sequences. In still another embodiment, a plasmid or vector of the invention contains at least one of the AAV-1 cap proteins of the invention. Most suitably, these AAV-1-derived vectors are assembled into viral vectors, as described herein.

A. AAV Viral Vectors

5

10

15

20

25

In one aspect, the present invention provides a recombinant AAV-1 viral vector produced using the AAV-1 capsid proteins of the invention. The packaged rAAV-1 virions of the invention may contain, in addition to a selected minigene, other AAV-1 sequences, or may contain sequences from other AAV serotypes.

Methods of generating rAAV virions are well known and the selection of a suitable method is not a limitation on the present invention. See, e.g., K. Fisher et al, J. Virol., 70:520-532 (1993) and US Patent 5,478,745. In one suitable method, a selected host cell is provided with the AAV sequence encoding a rep protein, the gene encoding the AAV cap protein and with the sequences for packaging and subsequent delivery. Desirably, the method utilizes the sequences encoding the AAV-1 rep and/or cap proteins of the invention.

In one embodiment, the rep/cap genes and the sequences for delivery are supplied by co-transfection of vectors carrying these genes and sequences. In one currently preferred embodiment, a cis (vector) plasmid, a trans plasmid containing the rep and cap genes, and a plasmid containing the adenovirus helper genes are co-transfected into a suitable cell line, e.g., 293. Alternatively, one or more of these functions may be provided in trans via separate vectors, or may be found in a suitably engineered packaging cell line.

An exemplary cis plasmid will contain, in 5' to 3' order, AAV 5' ITR, the selected transgene, and AAV 3' ITR. In one desirable embodiment, at least one of the AAV ITRs is a 143 nt AAV-1 ITR. However, other AAV serotype ITRs may be readily selected. Suitably, the full-length ITRs are utilized. However, one of skill in

.

5

10

15

20

25

30

12

the art can readily prepare modified AAV ITRs using conventional techniques.

Similarly, methods for construction of such plasmids is well known to those of skill in the art.

A trans plasmid for use in the production of the rAAV-1 virion particle may be prepared according to known techniques. In one desired embodiment, this plasmid contains the rep and cap proteins of AAV-1, or functional fragments thereof. Alternatively, the rep sequences may be from another selected AAV serotype.

The cis and trans plasmid may then be co-transfected with a wild-type helper virus (e.g., Ad2, Ad5, or a herpesvirus), or more desirably, a replication - defective adenovirus, into a selected host cell. Alternatively, the cis and trans plasmid may be co-transfected into a selected host cell together with a transfected plasmid which provides the necessary helper functions. Selection of a suitable host cell is well within the skill of those in the art and include such mammalian cells as 293 cells, HeLa cells, among others.

Alternatively, the cis plasmid and, optionally the trans plasmid, may be transfected into a packaging cell line which provides the remaining helper functions necessary for production of a rAAV containing the desired AAV-1 sequences of the invention. An example of a suitable packaging cell line, where an AAV-2 capsid is desired, is B-50, which stably expresses AAV-2 rep and cap genes under the control of a homologous P5 promoter. This cell line is characterized by integration into the cellular chromosome of multiple copies (at least 5 copies) of P5-rep-cap gene cassettes in a concatomer form. This B-50 cell line was deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209, on September 18, 1997 under Accession No. CRL-12401 pursuant to the provisions of the Budapest Treaty. However, the present invention is not limited as to the selection of the packaging cell line.

Exemplary transducing vectors based on AAV-1 capsid proteins have been tested both *in vivo and in vitro*, as described in more detail in Example 4. In these studies, it was demonstrated that recombinant AAV vector with an AAV-1 virion can transduce both mouse liver and muscle. These, and other AAV-1 based

13

gene therapy vectors which may be generated by one of skill in the art are beneficial for gene delivery to selected host cells and gene therapy patients since the neutralization antibodies of AAV-1 present in much of the human population exhibit different patterns from other AAV serotypes and therefore do not neutralize the AAV-1 virions. One of skill in the art may readily prepare other rAAV viral vectors containing the AAV-1 capsid proteins provided herein using a variety of techniques known to those of skill in the art. One may similarly prepare still other rAAV viral vectors containing AAV-1 sequence and AAV capsids of another serotype.

B. Other Viral Vectors

10

5

One of skill in the art will readily understand that the AAV-1 sequences of the invention can be readily adapted for use in these and other viral vector systems for *in vitro*, *ex vivo or in vivo* gene delivery. Particularly well suited for use in such viral vector systems are the AAV-1 ITR sequences, the AAV-1 rep, the AAV-1 cap, and the AAV-1 P5 promoter sequences.

15

20

25

30

For example, in one desirable embodiment, the AAV-1 ITR sequences of the invention may be used in an expression cassette which includes AAV-1 5' ITR, a non-AAV DNA sequences of interest (e.g., a minigene), and 3' ITR and which lacks functional rep/cap. Such a cassette containing an AAV-1 ITR may be located on a plasmid for subsequent transfection into a desired host cell, such as the cis plasmid described above. This expression cassette may further be provided with an AAV capsid of a selected serotype to permit infection of a cell or stably transfected into a desired host cell for packaging of rAAV virions. Such an expression cassette may be readily adapted for use in other viral systems, including adenovirus systems and lentivirus systems. Methods of producing Ad/AAV vectors are well known to those of skill in the art. One desirable method is described in PCT/US95/14018. However, the present invention is not limited to any particular method.

Another aspect of the present invention is the novel AAV-1 P5 promoter sequences which are located in the region spanning nt 236 - 299 of SEQ ID NO: 1. This promoter is useful in a variety of viral vectors for driving expression of a desired transgene.

10

15

20

25

Similarly, one of skill in the art can readily select other fragments of the AAV-1 genome of the invention for use in a variety of vector systems. Such vectors systems may include, e.g., lentiviruses, retroviruses, poxviruses, vaccinia viruses, and adenoviral systems, among others. Selection of these vector systems is not a limitation of the present invention.

C. Host Cells And Packaging Cell Lines

In yet another aspect, the present invention provides host cells which may be transiently transfected with AAV-1 nucleic acid sequences of the invention to permit expression of a desired transgene or production of a rAAV particle. For example, a selected host cell may be transfected with the AAV-1 P5 promoter sequences and/or the AAV-1 5' ITR sequences using conventional techniques. Providing AAV helper functions to the transfected cell lines of the invention results in packaging of the rAAV as infectious rAAV particles. Such cell lines may be produced in accordance with known techniques [see, e.g, US Patent No. 5,658,785], making use of the AAV-1 sequences of the invention.

Alternatively, host cells of the invention may be stably transfected with a rAAV expression cassette of the invention, and with copies of AAV-1 rep and cap genes. Suitable parental cell lines include mammalian cell lines and it may be desirable to select host cells from among non-simian mammalian cells. Examples of suitable parental cell lines include, without limitation, HeLa [ATCC CCL 2], A549 [ATCC Accession No. CCL 185], KB [CCL 17], Detroit [e.g., Detroit 510, CCL 72] and WI-38 [CCL 75] cells. These cell lines are all available from the American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 USA. Other suitable parent cell lines may be obtained from other sources and may be used to construct stable cell lines containing the P5 and/or AAV rep and cap sequences of the invention.

Recombinant vectors generated as described above are useful for delivery of the DNA of interest to cells.

15

III. METHODS OF DELIVERING GENES VIA AAV-1 DERIVED VECTORS

In another aspect, the present invention provides a method for delivery of a transgene to a host which involves transfecting or infecting a selected host cell with a recombinant viral vector generated with the AAV-1 sequences (or functional fragments thereof) of the invention. Methods for delivery are well known to those of skill in the art and are not a limitation of the present invention.

5

10

15

20

25

In one desirable embodiment, the invention provides a method for AAV-mediated delivery of a transgene to a host. This method involves transfecting or
infecting a selected host cell with a recombinant viral vector containing a selected
transgene under the control of sequences which direct expression thereof and AAV-1
capsid proteins.

Optionally, a sample from the host may be first assayed for the presence of antibodies to a selected AAV serotype. A variety of assay formats for detecting neutralizing antibodies are well known to those of skill in the art. The selection of such an assay is not a limitation of the present invention. See, e.g., Fisher et al, Nature Med., 3(3):306-312 (March 1997) and W. C. Manning et al, Human Gene Therapy, 9:477-485 (March 1, 1998). The results of this assay may be used to determine which AAV vector containing capsid proteins of a particular serotype are preferred for delivery, e.g., by the absence of neutralizing antibodies specific for that capsid serotype.

In one aspect of this method, the delivery of vector with AAV-1 capsid proteins may precede or follow delivery of a gene via a vector with a different serotype AAV capsid protein. Thus, gene delivery via rAAV vectors may be used for repeat gene delivery to a selected host cell. Desirably, subsequently administered rAAV vectors carry the same transgene as the first rAAV vector, but the subsequently administered vectors contain capsid proteins of serotypes which differ from the first vector. For example, if a first vector has AAV-2 capsid proteins, subsequently administered vectors may have capsid proteins selected from among the other serotypes, including AAV-1, AAV-3A, AAV-3B, AAV-4 and AAV-6.

16

Thus, a rAAV-1-derived recombinant viral vector of the invention provides an efficient gene transfer vehicle which can deliver a selected transgene to a selected host cell in vivo or ex vivo even where the organism has neutralizing antibodies to one or more AAV serotypes. These compositions are particularly well suited to gene delivery for therapeutic purposes. However, the compositions of the invention may also be useful in immunization. Further, the compositions of the invention may also be used for production of a desired gene product in vitro.

5

10

15

20

25

30

The above-described recombinant vectors may be delivered to host cells according to published methods. An AAV viral vector bearing the selected transgene may be administered to a patient, preferably suspended in a biologically compatible solution or pharmaceutically acceptable delivery vehicle. A suitable vehicle includes sterile saline. Other aqueous and non-aqueous isotonic sterile injection solutions and aqueous and non-aqueous sterile suspensions known to be pharmaceutically acceptable carriers and well known to those of skill in the art may be employed for this purpose.

The viral vectors are administered in sufficient amounts to transfect the cells and to provide sufficient levels of gene transfer and expression to provide a therapeutic benefit without undue adverse effects, or with medically acceptable physiological effects, which can be determined by those skilled in the medical arts. Conventional and pharmaceutically acceptable routes of administration include, but are not limited to, direct delivery to the liver, oral, intranasal, intravenous, intramuscular, subcutaneous, intradermal, and other parental routes of administration. Routes of administration may be combined, if desired.

Dosages of the viral vector will depend primarily on factors such as the condition being treated, the age, weight and health of the patient, and may thus vary among patients. For example, a therapeutically effective human dosage of the viral vector is generally in the range of from about 1 ml to about 100 ml of solution containing concentrations of from about 1 x 10⁹ to 1 x 10¹⁶ genomes virus vector. A preferred human dosage may be about 1 x 10¹³ to 1 x 10¹⁶ AAV genomes. The dosage will be adjusted to balance the therapeutic benefit against any side effects and

10

15

20

25

such dosages may vary depending upon the therapeutic application for which the recombinant vector is employed. The levels of expression of the transgene can be monitored to determine the frequency of dosage resulting in viral vectors, preferably AAV vectors containing the minigene. Optionally, dosage regimens similar to those described for therapeutic purposes may be utilized for immunization using the compositions of the invention. For *in vitro* production, a desired protein may be obtained from a desired culture following transfection of host cells with a rAAV containing the gene encoding the desired protein and culturing the cell culture under conditions which permits expression. The expressed protein may then be purified and isolated, as desired. Suitable techniques for transfection, cell culturing, purification, and isolation are known to those of skill in the art.

The following examples illustrate several aspects and embodiments of the invention.

Example 1 - Generation of Infectious Clone of AAV-1

The replicated form DNA of AAV-1 was extracted from 293 cells that were infected by AAV-1 and wild type adenovirus type 5.

A. Cell Culture and Virus

AAV-free 293 cells and 84-31 cells were provided by the human application laboratory of the University of Pennsylvania. These cells were cultured in Dulbecco's Modified Eagle Medium with 10% fetal bovine serum (Hyclone), penicillin (100 U/ml) and streptomycin at 37°C in a moisturized environment supplied with 5% C0₂. The 84-31 cell line constitutively expresses adenovirus genes E1a, Elb, E4/ORF6, and has been described previously [K. J. Fisher, J. Virol., 70:520-532 (1996)]. AAV-1 (ATCC VR-645) seed stock was purchased from American Type Culture Collection (ATCC, Manassas, VA). AAV viruses were propagated in 293 cells with wild type Ad5 as a helper virus.

B. Recombinant AAV Generation

The recombinant AAV viruses were generated by transfection using an adenovirus free method. Briefly, the cis plasmid (with AAV ITR), trans plasmid (with

18

AAV rep gene and cap gene) and helper plasmid (pFΔ13, with essential regions from the adenovirus genome) were simultaneously co-transfected into 293 cells in a ratio of 1:1:2 by calcium phosphate precipitation. The pFΔ13 helper plasmid has an 8 kb deletion in the adenovirus E2B region and has deletions in most of the late genes. This helper plasmid was generated by deleting the RsrII fragment from pFG140 (Microbix, Canada). Typically, 50 μg of DNA (cis:trans:PFΔ13 at ratios of 1:1:2, respectively) was transfected onto a 15 cm tissue culture dish. The cells were harvested 96 hours post-transfection, sonicated and treated with 0.5% sodium deoxycholate (37°C for 10 min). Cell lysates were then subjected to two rounds of a CsCl gradient. Peak fractions containing AAV vector were collected, pooled, and dialyzed against PBS before injecting into animals. To make rAAV virus with AAV-1 virion, the pAV1H or p5E18 (2/1) was used as the *trans* plasmid to provide rep and cap function.

5

10

For the generation of rAAV based on AAV-2, p5E18 was used as the trans plasmid since it greatly improved the rAAV yield. This plasmid, p5E18(2/2), 15 expresses AAV-2 Rep and Cap and contains a P5 promoter relocated to a position 3' to the Cap gene, thereby minimizing expression of Rep78 and Rep68. The strategy was initially described by Li et al, <u>J. Virol.</u>, 71:5236-5243 (1997). P5E18(2/2) was constructed in the following way. The previously described pMMTV-trans vector 20 (i.e., the mouse mammary tumor virus promoter substituted for the P5 promoter in an AAV-2-based vector) was digested with SmaI and ClaI, filled in with the Klenow enzyme, and then recircularized with DNA ligase. The resulting construct was digested with XbaI, filled in, and ligated to the blunt-ended BamHI-XbaI fragment from pCR-p5, constructed in the following way. The P5 promoter of AAV was 25 amplified by PCR and the amplified fragment was subsequently cloned into pCR2.1 (Invitrogen) to yield pCR-P5. The helper plasmid pAV1H was constructed by cloning the BfaI fragment of pAAV-2 into pBluescript II-SK(+) at the BcorV and SmaI sites. The 3.0-kb XbaI-KpnI fragment from p5E18(2/2), the 2.3-kb XbaI-KpnI fragment from pAV1H, and the 1.7-kb KpnI fragment from p5E18(2/2) were incorporated into 30 a separate plasmid P5E18(2/1), which contains AAV-2 Rep, AAV-1 Cap, and the

19

AAV-2 P5 promoter located 3' to the Cap gene. Plasmid p5E18(2/1) produced 10- to 20-fold higher quantities of the vector than pAV1H (i.e., 10¹² genomes/50 15-cm² plates).

C. <u>DNA Techniques</u>

5

10

15

30

Hirt DNA extraction was performed as described in the art with minor modification [R.J. Samulski et al., Cell, 33:135-143 (1983)]. More particularly, Hirst solution without SDS was used instead of using original Hirt solution containing SDS. The amount of SDS present in the original Hirst solution was added after the cells had been fully suspended. To construct AAV-1 infectious clone, the Hirt DNA from AAV-1 infected 293 cells was repaired with Klenow enzyme (New England Biolabs) to ensure the ends were blunt. The treated AAV-1 Hirt DNA was then digested with BamHI and cloned into three vectors, respectively. The internal BamHI was cloned into pBlueScript II-SK+ cut with BamHI to get pAV1-BM. The left and right fragments were cloned into pBlueScript II-SK+ cut with BamHI + EcoRV to obtain pAV1-BL and pAV1-BR, respectively. The AAV sequence in these three plasmids were subsequently assembled into the same vector to get AAV-1 infectious clone pAAV-1. The helper plasmid for recombinant AAV-1 virus generation was constructed by cloning the Bfa I fragment of pAAV-1 into pBlueScript II-SK+ at the EcoRV site.

Analysis of the Hirt DNA revealed three bands, a dimer at 9.4 kb, a monomer at 4.7 kb and single-stranded DNA at 1.7 kb, which correlated to different replication forms of AAV-1. The monomer band was excised from the gel and then digested with BamHI. This resulted in three fragments of 1.1 kb, 0.8 kb and 2.8 kb. This pattern is in accordance with the description by Bantel-schaal and zur Hausen,

Virol., 134(1):52-63 (1984). The 1.1 kb and 2.8 kb BamHI fragments were cloned into pBlueScript-KS(+) at BamHI and EcoRV site. The internal 0.8 kb fragment was cloned into BamHI site of pBlueScript-KS(+).

These three fragments were then subcloned into the same construct to obtain a plasmid (pAAV-1) that contained the full sequence of AAV-1. The pAAV-1 was then tested for its ability to rescue from the plasmid backbone and package

20

infectious virus. The pAAV-1 was then transfected to 293 cells and supplied with adenovirus type as helper at MOI 10. The virus supernatant was used to reinfect 293 cells.

For Southern blot analysis, Hirt DNA was digested with *DpnI* to remove bacteria-borne plasmid and probed with internal *BamHI* fragment of AAV-1. The membrane was then washed at high stringency conditions, which included: twice 30 minutes with 2X SSC, 0.1% SDS at 65°C and twice 30 minutes with 0.1X SSC, 0.1% SDS at 65°C. The membrane was then analyzed by both phosphor image and X-ray autoradiography. The results confirmed that pAAV-1 is indeed an infectious clone of AAV serotype 1.

Example 2 - Sequencing Analysis of AAV-1

5

10

15

20

25

The entire AAV-1 genome was then determined by automatic sequencing and was found to be 4718 nucleotides in length (Figs. 1A-1C). For sequencing, an ABI 373 automatic sequencer as used to determine the sequences for all plasmids and PCR fragments related to this study using the FS dye chemistry. All sequences were confirmed by sequencing both plus and minus strands. These sequences were also confirmed by sequencing two independent clones of pAV-BM, pAV-BL and pAV-BR. Since the replicated form of AAV-1 DNA served as the template for sequence determination, these sequences were also confirmed by sequencing a series of PCR products using original AAV-1 seed stock as a template.

The length of AAV-1 was found to be within the range of the other serotypes: AAV-3 (4726 nucleotides), AAV-4 (4774 nucleotides), AAV-2 (4681 nucleotides), and AAV-6 (4683 nucleotides).

The AAV-1 genome exhibited similarities to other serotypes of adenoassociated viruses. Overall, it shares more than 80% identity with other known AAV
viruses as determined by the computer program Megalign using default settings
[DNASTAR, Madison, WI]. The key features in AAV-2 can also be found in AAV1. First, AAV-1 has the same type of inverted terminal repeat which is capable of
forming T-shaped hairpin structures, despite the differences at the nucleotide level

21

(Figs. 2 and 3). The sequences of right ITRs and left ITRs of AAV-1 are identical. The AAV TR sequence is subdivided into A, A', B, B', C, C', D and D' [Bern, cited above].

These AAV ITR sequences are also virtually the same as those found in AAV-6 right ITR, there being one nucleotide difference in each of A and A' sequence, and the last nucleotide of the D sequence. Second, the AAV-2 rep binding motif [GCTCGCTCGCTG (SEQ ID NO: 20)] is well conserved. Such motif can also be found in the human chromosome 19 AAV-2 pre-integration region. Finally, non-structural and structural coding regions, and regulatory elements similar to those of other AAV serotypes also exist in AAV-1 genome.

5

10

15

20

25

30

Although the overall features of AAV terminal repeats are very much conserved, the total length of the AAV terminal repeat exhibits divergence. The terminal repeat of AAV-1 consists of 143 nucleotides while those of AAV-2, AAV-3, and AAV-4 are about 145 or 146 nucleotides. The loop region of AAV-1 ITR most closely resembles that of AAV-4 in that it also uses TCT instead of the TTT found in AAV-2 and AAV-3. The possibility of sequencing error was eliminated using restriction enzyme digestion, since these three nucleotides are part of the SacI site (gagete; nt 69-74 of SEQ ID NO: 1). The p5 promoter region of AAV-1 shows more variations in nucleotide sequences with other AAV serotypes. However, it still maintains the critical regulatory elements. The two copies of YY1 [See, Fig. 1A-1C] sites seemed to be preserved in all known AAV serotypes, which have been shown to be involved in regulating AAV gene expression. In AAV-4, there are 56 additional nucleotides inserted between YY1 and E-box/USF site, while in AAV-1, there are 26 additional nucleotides inserted before the E-box/USF site. The p19 promoter, p40 promoter and polyA can also be identified from the AAV-1 genome by analogy to known AAV serotypes, which are also highly conserved.

Thus, the analysis of AAV terminal repeats of various serotypes showed that the A and A' sequence is very much conserved. One of the reasons may be the Rep binding motif (GCTC)₃GCTG [SEQ ID NO: 20]. These sequences appear to be essential for AAV DNA replication and site-specific integration. The same sequence

10

15

20

25

has also been shown to be preserved in a monkey genome [Samulski, personal communication]. The first 8 nucleotides of the D sequence are also identical in all known AAV serotypes. This is in accordance with the observation of the Srivastava group that only the first 10 nucleotides are essential for AAV packaging [X.S. Wang et al, J. Virol., 71:3077-3082 (1997); X.S. Wang et al, J. Virol., 71:1140-1146 (1997)]. The function of the rest of the D sequences still remain unclear. They may be somehow related to their tissue specificities. The variation of nucleotide in B and C sequence may also suggest that the secondary structure of the ITRs is more critical for its biological function, which has been demonstrated in many previous publications.

Example 3 - Comparison of AAV-1 Sequences

The nucleotide sequences of AAV-1, obtained as described above, were compared with known AAV sequences, including AAV-2, AAV-4 and AAV-6 using DNA Star Megalign. This comparison revealed a stretch of 71 identical nucleotides shared by AAV-1, AAV-2 and AAV-6. See, Figs. 1A-1C.

This comparison further suggested that AAV-6 is a hybrid formed by homologous recombination of AAV-1 and AAV-2. See, Figs. 3A and 3B. These nucleotides divide the AAV-6 genome into two regions. The 5' half of AAV-6 of 522 nucleotides is identical to that of AAV-2 except in 2 positions. The 3' half of AAV-6 including the majority of the rep gene, complete cap gene and 3' ITR is 98% identical to AAV-1.

Biologically, such recombination may enable AAV-1 to acquire the ability to transmit through the human population. It is also interesting to note that the ITRs of AAV-6 comprise one AAV-1 ITR and one AAV-2 ITR. The replication model of defective parvovirus can maintain this special arrangement. Studies on AAV integration have shown that a majority of AAV integrants carries deletions in at least one of the terminal repeats. These deletions have been shown to be able to be repaired through gene conversion using the other intact terminal repeat as a template. Therefore, it would be very difficult to maintain AAV-6 as a homogenous population

10

15

20

25

when an integrated copy of AAV-6 is rescued from host cells with helper virus infection. The AAV-6 with two identical AAV-2 ITRs or two identical AAV-1 ITRs should be the dominant variants. The AAV-6 with two AAV-1 ITRs has been observed by Russell's group [Rutledge, cited above (1998)]. So far there is no report on AAV-6 with two AAV-2 ITRs. Acquirement of AAV-2 P5 promoter by AAV-6 may have explained that AAV-6 have been isolated from human origin while AAV-1 with the same virion has not. The regulation of P5 promoter between different species of AAV may be different *in vivo*. This observation suggests the capsid proteins of AAV were not the only determinants for tissue specificity.

Although it is clear that AAV-6 is a hybrid of AAV-1 and AAV-2, AAV-6 has already exhibited divergence from either AAV-1 or AAV-2. There are two nucleotide differences between AAV-6 and AAV-2 in their first 450 nucleotides. There are about 1% differences between AAV-6 and AAV-1 in nucleotide levels from nucleotides 522 to the 3' end. There also exists a quite divergent region (nucleotide 4486-4593) between AAV-6 and AAV-1 (Figs. 1A-1C). This region does not encode any known proteins for AAVs. These differences in nucleotide sequences may suggest that AAV-6 and AAV-1 have gone through some evolution since the recombination took place. Another possible explanation is that there exists another variant of AAV-1 which has yet to be identified. So far, there is no evidence to rule out either possibility. It is still unknown if other hybrids (AAV-2 to AAV-4, etc.) existed in nature.

The coding region of AAV-1 was deduced by comparison with other known AAV serotypes. Table 1 illustrates the coding region differences between AAV-1 and AAV-6. The amino acid residues are deduced according to AAV-2.

With reference to the amino acid position of AAV-1, Table 1 lists the amino acids of AAV-1 which have been changed to the corresponding ones of AAV-6. The amino acids of AAV-1 are shown to the left of the arrow. Reference may be made to SEQ ID NO: 5 of the amino acid sequence of AAV-1 Rep 78 and to SEQ ID NO: 13 for the amino acid sequence of AAV-1 VP1.

Table 1

Coding region variations between AAV-1 and AAV-6

24

Rep prote	ein (Rep78)		Cap pro	tein (VP1)
Position(s)	Amino acids		Position(s)	Amino acids
28	S→N		129	L→F
191	Q-H		418	E→D
192	H-D		531	E∹K
308	E-D		584	F→L
]	598	A-V
			642	N−H

It was surprising to see that the sequence of the AAV-1 coding region is
almost identical to that of AAV-6 from position 452 to the end of coding region
(99%). The first 508 nucleotides of AAV-6 have been shown to be identical to those
of AAV-2 [Rutledge, cited above (1998)]. Since the components of AAV-6 genome
seemed to be AAV-2 left ITR - AAV-2 p5 promoter - AAV-1 coding region - AAV1 right ITR, it was concluded that AAV-6 is a naturally occurred hybrid between
AAV-1 and AAV-2.

Example 4 - Gene Therapy Vector Based on AAV-1

5

20

Recombinant gene transfer vectors based on AAV-1 viruses were constructed by the methods described in Example 1. To produce a hybrid recombinant virus with AAV-1 virion and AAV-2 ITR, the AAV-1 trans plasmid (pAV1H) and the AAV-2 cis-lacZ plasmid (with AAV-2 ITR) were used. The AAV-2 ITR was used in this vector in view of its known ability to direct site-specific integration. Also constructed for use in this experiment was an AAV-1 vector carrying the green fluorescent protein (GFP) marker gene under the control of the immediate early promoter of CMV using pAV1H as the trans plasmid.

10

15

20

25

A. <u>rAAV-1 Viruses Transfect Host Cells in Vitro</u>

84-31 cells, which are subclones of 293 cells (which express adenovirus E1a, E1b) which stably express E4/ORF5, were infected with rAAV-1 GFP or rAAV-lacZ. High levels of expression of GFP and lacZ was detected in the cultured 84-31 cells. This suggested that rAAV-1 based vector was very similar to AAV-2 based vectors in ability to infect and expression levels.

B. <u>rAAV-1 Viruses Transfect Cells in Vivo</u>

The performance of AAV-1 based vectors was also tested *in vivo*. The rAAV-1 CMV-α1AT virus was constructed as follows. The EcoRI fragment of pAT85 (ATCC) containing human α1-antitrypsin (α1AT) cDNA fragment was blunted and cloned into PCR (Promega) at a Small site to obtain PCR-α1AT. The CMV promoter was cloned into PCR-α1AT at the Xball site. The Alb-α1AT expression cassette was removed by XhoI and ClaI and cloned into pAV1H at the Xball site. This vector plasmid was used to generate AAV-1-CMV-α1AT virus used in the experiment described below.

For screening human antibodies against AAV, purified AAV virus is lysed with Ripa buffer (10 mM Tris pH 8.2, 1% Triton X-100, 1% SDS, 0.15 M NaCl) and separated in 10% SDS-PAGE gel. The heat inactivated human serum was used at a 1 to 1000 dilution in this assay. The rAAV-1 CMV- α 1AT viruses were injected into Rag-1 mice through tail vein injection at different dosages. The concentration of human α 1-antitrypsin in mouse serum was measured using ELISA. The coating antibody is rabbit anti-human human α 1-antitrypsin (Sigma). The goatantihuman α 1-antitrypsin (Sigma) was used as the primary detection antibodies. The sensitivity of this assay is around 0.3 ng/ml to 30 ng/ml. The expression of human α 1-antitrypsin in mouse blood can be detected in a very encouraging level. This result is shown in Table 2.

26

Table 2
Human Antitrypsin Expressed in Mouse Liver

Amount of virus injected	Week 2 (ng/ml)	Week 4 (ng/ml)
2x10 ¹⁰ genomes	214.2	171.4
1x10 ¹⁰ genomes	117.8	109.8
5x10 ¹⁰ genomes	64.5	67.8
2.5x10 ¹⁰ genomes	30.9	58.4

rAAV-1 CMV-lacZ viruses were also injected into the muscle of
C57BL6 mice and similar results were obtained. Collectively, these results suggested
that AAV-1 based vector would be appropriate for both liver and muscle gene
delivery.

Example 5 - Neutralizing Antibodies Against AAV-1

Simple and quantitative assays for neutralizing antibodies (NAB) to AAV-1 and AAV-2 were developed with recombinant vectors. A total of 33 rhesus monkeys and 77 normal human subjects were screened.

A. Nonhuman Primates

5

15

20

25

Wild-caught juvenile rhesus monkeys were purchased from Covance (Alice, Tex.) and LABS of Virginia (Yemassee, SC) and kept in full quarantine. The monkeys weighed approximately 3 to 4 kg. The nonhuman primates used in the Institute for Human Gene Therapy research program are purposefully bred in the United States from specific-pathogen-free closed colonies. All vendors are US Department of Agriculture class A dealers. The rhesus macaques are therefore not infected with important simian pathogens, including the tuberculosis agent, major simian lentiviruses (simian immunodeficiency virus and simian retroviruses), and cercopithecine herpesvirus. The animals are also free of internal and external parasites. The excellent health status of these premium animals minimized the potential for extraneous variables. For this study, serum was obtained from monkeys prior to initiation of any protocol.

27

NAB titers were analyzed by assessing the ability of serum antibody to inhibit the transduction of reporter virus expressing green fluorescent protein (GFP) (AAV1-GFP or AAV2-GFP) into 84-31 cells. Various dilutions of antibodies preincubated with reporter virus for 1 hour at 37°C were added to 90% confluent cell cultures. Cells were incubated for 48 hours and the expression of green fluorescent protein was measured by FluoroImaging (Molecular Dynamics). NAB titers were calculated as the highest dilution at which 50% of the cells stained green.

Analysis of NAB in rhesus monkeys showed that 61% of animals tested positive for AAV-1; a minority (24%) has NAB to AAV-2. Over one-third of animals had antibodies to AAV-1 but not AAV-2 (i.e., were monospecific for AAV-1), whereas no animals were positive for AAV-2 without reacting to AAV-1. These data support the hypothesis that AAV-1 is endemic in rhesus monkeys. The presence of true AAV-2 infections in this group of nonhuman primates is less clear, since cross-neutralizing activity of an AAV-1 response to AAV-2 can not be ruled out. It is interesting that there is a linear relationship between AAV-2 NAB and AAV-1 NAB in animals that had both.

B. Humans

5

10

15

20

25

For these neutralization antibody assays, human serum samples were incubated at 56°C for 30 min to inactivate complement and then diluted in DMEM. The virus (rAAV or rAd with either lacZ or GFP) was then mixed with each serum dilution (20X, 400X, 2000X, 4000X, etc.) and incubated for 1 hour at 37°C before applied to 90% confluent cultures of 84-31 cells (for AAV) or Hela cells (for adenovirus) in 96-well plates. After 60 minutes of incubation at culture condition, 100 µl additional media containing 20% FCS was added to make final culture media containing 10% FCS.

> 28 The result is summarized in Table 3.

Table 3

Adenovirus	AAV-1	AAV-2	# of samples	Percentage
-	-	-	41	53.2%
+	_		16	20.8%
-	+	- ,	0	0.0%
•	-	+	2	2.6%
-	+	+	2	2.6%
+	-	+	3	3.9%
+	+	-	0	0.0%
+	+	+	13	16.9%
		Total	77	100%

5

The human neutralizing antibodies against these three viruses seemed to be unrelated since the existence of neutralizing antibodies against AAV are not indications for antibodies against adenovirus. However, AAV requires adenovirus as helper virus, in most of the cases, the neutralizing antibodies against AAV correlated with the existence of neutralizing antibodies to adenovirus. Among the 77 human serum samples screened, 41% of the samples can neutralize the infectivity of recombinant adenovirus based on Ad5. 15/77 (19%) of serum samples can neutralize the transduction of rAAV-1 while 20/77 (20%) of the samples inhibit rAAV-2 transduction at 1 to 80 dilutions or higher. All serum samples positive in neutralizing antibodies for AAV-1 in are also positive for AAV-2. However, there are five (6%) rAAV-2 positive samples that failed to neutralize rAAV-1. In samples that are positive for neutralizing antibodies, the titer of antibodies also varied in the positive ones. The results from screening human sera for antibodies against AAVs supported the conclusion that AAV-1 presents the same epitome as that of AAV-2 to interact

10

15

25

20

29

with cellular receptors since AAV-1 neutralizing human serums can also decrease the infectivity of AAV-2. However, the profile of neutralizing antibodies for these AAVs is not identical, there are additional specific receptors for each AAV serotype.

Example 6 - Recombinant AAV Viruses Exhibit Tissue Tropism

5

10

15

20

25

The recombinant AAV-1 vectors of the invention and the recombinant AAV-2 vectors [containing the gene encoding human α 1-antitrypsin (α 1AT) or murine erythropoietin (Epo) from a cytomegalovirus-enhanced β -actin promoter (CB)] were evaluated in a direct comparison to equivalent copies of AAV-2 vectors containing the same vector genes.

Recombinant viruses with AAV-1 capsids were constructed using the techniques in Example 1. To make rAAV with AAV-1 virions, pAV1H or p5E18 (2/1) was used as the *trans* plasmid to provide Rep and Cap functions. For the generation of the rAAV based on AAV-2, p5E18(2/2) was used as the *trans* plasmid, since it greatly improved the rAAV yield. [Early experiments indicated similar *in vivo* performances of AAV-1 vectors produced with pAV1H and p5E19 (2/1). All subsequent studies used AAV-1 vectors derived from p5E18(2/1) because of the increased yield.]

Equivalent stocks of the AAV-1 and AAV-2 vectors were injected intramuscularly (5 x 10¹⁰ genomes) or liver via the portal circulation (1 x 10¹¹ genomes) into immunodeficient mice, and the animals (four groups) were analyzed on day 30 for expression of transgene. See, Figs. 4A and 4B.

AAV-2 vectors consistently produced 10- to 50-fold more serum erythropoietin or α 1-antitrypsin when injected into liver compared to muscle. (However, the AAV-1-delivered genes did achieve acceptable expression levels in the liver.) This result was very different from that for AAV-1 vectors, with which muscle expression was equivalent to or greater than liver expression. In fact, AAV-1 outperformed AAV-2 in muscle when equivalent titers based on genomes were administered.

30

Example 7 - Gene Delivery via rAAV-1

5

10

15

20

25

30

C57BL/6 mice (6- to 8-week old males, Jackson Laboratories) were analyzed for AAV mediated gene transfer to liver following intrasplenic injection of vector (i.e., targeted to liver). A total of 10¹¹ genome equivalents of rAAV-1 or rAAV-2 vector were injected into the circulation in 100 μl buffered saline. The first vector contained either an AAV-1 capsid or an AAV-2 capsid and expressed α1AT under the control of the chicken β-actin (CB) promoter. Day 28 sera were analyzed for antibodies against AAV-1 or AAV-2 and serum α1AT levels were checked. Animals were then injected with an AAV-1 or AAV-2 construct expressing erythropoietin (Epo, also under the control of the CB promoter). One month later sera was analyzed for serum levels of Epo. The following groups were analyzed (Figs. 5A-5D).

In Group 1, vector 1 was AAV-2 expressing α 1AT and vector 2 was AAV-2 expressing Epo. Animals generated antibodies against AAV-2 following the first vector administration which prevented the readministration of the AAV-2 based vector. There was no evidence for cross-neutralizing the antibody to AAV-1.

In Group 2, vector 1 was AAV-1 expressing α 1AT while vector 2 was AAV-1 expressing Epo. The first vector administration did result in significant α 1AT expression at one month associated with antibodies to neutralizing antibodies to AAV-1. The animals were not successfully readministered with the AAV-1 Epo expressing construct.

In Group 3, the effectiveness of an AAV-2 vector expressing Epo injected into a naive animal was measured. The animals were injected with PBS and injected with AAV-2 Epo vector at day 28 and analyzed for Epo expression one month later. The neutralizing antibodies were evaluated at day 28 so we did not expect to see anything since they received PBS with the first vector injection. This shows that in naive animals AAV-2 is very efficient at transferring the Epo gene as demonstrated by high level of serum Epo one month later.

Group 4 was an experiment similar to Group 3 in which the animals originally received PBS for vector 1 and then the AAV-1 expressing Epo construct 28 days later. At the time of vector injection, there obviously were no antibodies to either

10

15

20

AAV-1 or AAV-2. The AAV-1 based vector was capable of generating significant expression of Epo when measured one month later.

Group 5 is a cross-over experiment where the initial vector is AAV-2 expressing α1AT followed by the AAV-1 construct expressing Epo. The animals, as expected, were efficiently infected with the AAV-2 vector expressing α1AT as shown by high levels of the protein in blood at 28 days. This was associated with significant neutralizing antibodies to AAV-2. Importantly, the animals were successfully administered AAV-1 following the AAV-2 vector as shown by the presence of Epo in serum 28 days following the second vector administration. At the time of this vector administration, there was high level AAV-2 neutralizing antibodies and very low cross-reaction to AAV-1. The level of Epo was slightly diminished possibly due to a small amount of cross-reactivity. Group 6 was the opposite cross-over experiment in which the initial vector was AAV-1 based, whereas the second experiment was AAV-2 based. The AAV-1 vector did lead to significant gene expression of α1AT, which also resulted in high level AAV-1 neutralizing antibody. The animals were very efficiently administered AAV-2 following the initial AAV-1 vector as evidenced by high level Epo.

A substantially identical experiment was performed in muscle in which 5×10^{10} genomes were injected into the tibialis anterior of C57BL/6 mice as a model for muscle directed gene therapy. The results are illustrated in Figs. 6A-6D and are essentially the same as for liver.

In summary, this experiment demonstrates the utility of using an AAV-1 vector in patients who have pre-existing antibodies to AAV-2 or who had initially received an AAV-2 vector and need readministration.

25 Example 8 - Construction of Recombinant Viruses Containing AAV-1 ITRs

This example illustrates the construction of recombinant AAV vectors which contain AAV-1 ITRs of the invention.

An AAV-1 cis plasmid is constructed as follows. A 160 bp Xho-NruI AAV-1 fragment containing the AAV-1 5' ITR is obtained from pAV1-BL. pAV1-BL was

32

generated as described in Example 1. The Xho-NruI fragment is then cloned into a second pAV1-BL plasmid at an XbaI site to provide the plasmid with two AAV-1 ITRs. The desired transgene is then cloned into the modified pAV-1BL at the NruI and BamHI site, which is located between the AAV-1 ITR sequences. The resulting AAV-1 cis plasmid contains AAV-1 ITRs flanking the transgene and lacks functional AAV-1 rep and cap.

5

10

15

20

Recombinant AAV is produced by simultaneously transfecting three plasmids into 293 cells. These include the AAV-1 cis plasmid described above; a trans plasmid which provides AAV rep/cap functions and lacks AAV ITRs; and a plasmid providing adenovirus helper functions. The rep and/or cap functions may be provided in trans by AAV-1 or another AAV serotype, depending on the immunity profile of the intended recipient. Alternatively, the rep or cap functions may be provided in cis by AAV-1 or another serotype, again depending on the patient's immunity profile.

In a typical cotransfection, 50 µg of DNA (cis:trans:helper at ratios of 1:1:2, respectively) is transfected onto a 15 cm tissue culture dish. Cells are harvested 96 hours post transfection, sonicated and treated with 0.5% sodium deoxycholate (37° for 10 min). Cell lysates are then subjected to 2-3 rounds of ultracentrifugation in a cesium gradient. Peak fractions containing rAAV are collected, pooled and dialyzed against PBS. A typical yield is 1 x 10¹³ genomes/10⁹ cells.

Using this method, one recombinant virus construct is prepared which contains the AAV-1 ITRs flanking the transgene, with an AAV-1 capsid. Another recombinant virus construct is prepared with contains the AAV-1 ITRs flanking the transgene, with an AAV-2 capsid.

All publications cited in this specification are incorporated herein by reference.

While the invention has been described with reference to a particularly preferred embodiments, it will be appreciated that modifications can be made without departing from the spirit of the invention. Such modifications are intended to fall within the scope of the claims.

What is claimed is:

- 1. An isolated AAV-1 nucleic acid molecule comprising a sequence selected from the group consisting of:
 - (a) SEQ ID NO: 1;
 - (b) a DNA sequence complementary to SEQ ID NO: 1;
 - (c) cDNA complementary to (a) or (b); and
 - (d) RNA complementary to any of (a) to (c).
- 2. A nucleic acid molecule comprising an AAV-1 inverted terminal repeat (ITR) sequence selected from the group consisting of:
 - (a) nt 1 to 143 of SEQ ID NO: 1;
 - (b) nt 4576 to 4718 of SEQ ID NO: 1;
 - (c) a nucleic acid sequence complementary to (a) or (b); and
 - (d) a functional fragment of (a), (b), or (c).
- 3. A recombinant vector comprising a 5' AAV-1 inverted terminal repeat (ITR) and a selected transgene, wherein said ITR has the sequence selected from the group consisting of:
 - (a) nt 1 to 143 of SEQ ID NO: 1;
 - (b) a nucleic acid sequence complementary to (a); and
 - (c) a functional fragment of (a) or (b).
- 4. The recombinant vector according to claim 3, wherein said vector further comprises a 3' AAV-1 ITR.

- 5. A recombinant vector comprising a 3' AAV-1 inverted terminal repeat (ITR) and a selected transgene, wherein said ITR has the sequence selected from the group consisting of:
 - (a) nt 4576 to 4718 of SEQ ID NO: 1;
 - (b) a nucleic acid sequence complementary to (a); and
 - (c) a functional fragment of (a) or (b).
- 6. The recombinant vector according to claim 5, wherein said vector further comprises a 5' AAV-1 ITR.
- 7. The recombinant vector according to any of claims 3-6, wherein said vector further comprises AAV-1 capsid proteins having the sequence of SEQ ID NO: 13, 15 or 17 or functional fragments thereof.
- 8. The recombinant vector according to any of claims 3-6, wherein said vector further comprises adenovirus sequences.
- 9. A recombinant vector comprising an AAV-1 P5 promoter having the sequence of nt 236 to 299 of SEQ ID NO: 1 or a functional fragment thereof.
- 10. A nucleic acid molecule encoding AAV-1 helper functions, said molecule comprising an AAV rep coding region and an AAV cap coding region, wherein said cap coding region comprises at least one member is selected from the group consisting of:
 - (a) vp1, nt 2223 to 4431 of SEQ ID NO: 1;
 - (b) vp2, nt 2634 to 4432 of SEQ ID NO: 1; and
 - (c) vp3, nt 2829 to 4432 of SEQ ID NO: 1.

- 11. A nucleic acid molecule encoding AAV-1 helper functions, said molecule comprising an AAV rep coding region and an AAV cap coding region, wherein said rep coding region comprises an AAV-1 rep coding region comprising at least one member selected from the group consisting of:
 - (a) rep 78, nt 335 to 2304 of SEQ ID NO: 1;
- (b) rep 68, nt 335 to 2272 of SEQ ID NO: 1 or the cDNA corresponding thereto;
 - (c) rep 52, nt 1007 to 2304 of SEQ ID NO: 1; and
- (d) rep 40, nt 1007 to 2272 of SEQ ID NO: 1 or the cDNA corresponding thereto.
- 12. A host cell transduced with a recombinant viral vector according to any of claims 3-6.
- 13. A host cell transduced with a nucleic acid molecule according to any of claims 1, 2, 10 or 11.
- 14. A host cell stably transduced with an AAV-1 P5 promoter having the sequence of nt 236 to 299 of SEQ ID NO: 1.
- 15. A pharmaceutical composition comprising a carrier and a virus comprising the vector according to any of claims 3-6.
- 16. A pharmaceutical composition comprising a carrier and a virus comprising the vector according to claim 7.
- 17. A pharmaceutical-composition comprising a carrier and a virus comprising the vector according to claim 8.

- 18. A method for AAV-mediated delivery of a transgene comprising the step of delivering to a host cell an AAV virion which comprises:
- (a) a capsid comprising at least one capsid protein encoded by an AAV-1 cap gene; and
- (b) a DNA molecule comprising a transgene under the control of regulatory sequences directing its expression.
- 19. A method for AAV-mediated delivery of a transgene to a host comprising the steps of:
- (a) assaying a sample from the host to determine the presence of neutralizing antibodies specific against any serotype of AAV; and
 - (b) delivering to the host an AAV virion which comprises:
- (i) a capsid comprising at least one capsid protein encoded by a cap gene of an AAV serotype against which the host has no antibodies as determined in step (a); and
- (ii) a DNA molecule comprising a transgene under the control of regulatory sequences directing its expression.
- 20. The method according to claim 19, comprising the additional step of repeating steps (a) and (b).
- 21. Use of an AAV virion which comprises a capsid comprising (a) at least one capsid protein encoded by a cap gene of an AAV serotype against which the host has antibodies, and (b) a DNA molecule comprising a transgene operably linked to regulatory sequences directing its expression,

in the preparation of a medicament for delivery of a transgene to a host, wherein said host has no preexisting neutralizing antibodies against the AAV serotype of said cap gene.

- A method for delivery of a transgene comprising the step of delivering to a host cell a recombinant virus comprising a recombinant vector according to any of claims 3-8.
- 23. A method for producing a selected gene product comprising the steps of transfecting a mammalian cell with the molecule according to claim 1 or a functional fragment thereof and culturing said cell under conditions suitable to express said gene product.

FIG IA

	Rep binding site	
AAV-1	1 ttgcccactcccttctgcgggttggctcggtggggcctggggaccaaaggtcggcagaaggggagaggcgctctgctgctgctgggggcgcgagggggggg	, 120 120
AAV-	6gtg.g	. 120
AAV-1	TRS E box/USF 1 presente est est est est est est est est est e	237
AAV-1	2 .e	. 322
****		•••
	YYI PS/TATA YYI/p5 RNA Rep78/68	
AAV-1	CACATTTTCCGACACCACCTGCCCATTTAGGCTATATAGGCCGAGTCAGCGAGCAGCATTTCGAC-CGCGAAATTTGAACGAGCAGCCATGCCGGGCTTCTACGAGATCC) 356 342
AAV-6		341
AAV-1	L TEATCAAGGTGCCGAGCGACCTGCACGACCACCTGCCCGGCATTTCTGACTCGGTTTGTGAGCTGGGTGGCCCACAAGGAATGGGGGCTGCCCCGGGATTCTGACATGGATCTGAATCTGA	476
YYA-3	3TCCTGTCAGCATGATTGATTGATTGATTGATTGATTGATTGATTGATTGATTGATTGATTGATTGATT	462
AAV-6	,	481
	· · · · · · · · · · · · · · · · · · ·	• •
AAV-1	TTOMOCHOGOCHOCCCTCACCCGTCCCACGAAGCTGCCGCCCCACTTCCTCGTCCAATGGCCCCCGCGTGAGTAAGGCCCCGCGAGGCCCTCTTCTTTGTTCAGTACGAAGGGCGAGTCCT	596
AAV-6		581
AAV-1	. ACTTCCACCTCCATATTCTGGGGGGCCACGGGGCCAAATCCATGGTGCTGGGCCGGTTCCTGAGTCAGATTAGGGACAAGCTGGTGCAGACCATCTACGGGGGATGGAGCCGACCA	716
117-2		702
774-9		701
AAV-1	TOCCCAACTGGTTGGCGGTGAGCAAGAGGGGGTAATGGGGGGGG	#36
77A-1	A	822
	•	
	P19/TATA P19 RNA	
AAV-1	T. AC. T. T. C. G. T. CA. G. T. T. T. G.	942
AAV-6	TACTCCGTCA.GTT.GTGGTCGGAAA.	941
	Rep52/40	
44V-1	ACCCCANTICTORCGCCCTGTCATCCCGGTCAAAAACCTCCGCCGCCTACATCCACCTGGTCGGTGGGTG	1076
MA-3	.T	1062
A.V.		1001
	CCTCGTACATCTCCTTCAACGCGGCTTCCAACTCGGGGGTCCCAGATCAAGGCCGGTTCTGGACAATGCCGGCAAGATCATGGCGGTGACCAAATCCGGGGCCCGACTACCTGGTAGGCCCCG	
MV-3	T.G. C. A. T.CT. G. A. T. AGC. T. A. C. G. A. G. AGC.	1182
WA-1		1181
•	•	
WA-1	CTCCGCCCGGGCATTAAAACCAACCGCATCTACCGCATCCTGGGGCTGAACCGCATGAACCTGCCTG	1316
W∆-3	AGCOTG.ATCC.GTGTTAAATTA.A	1302
WA-0		1301
WA-1	ACACCATCTGGCCTGTTTGGGCCGGCCACCACGGGCAAGACCAACACGCGGAAGCCATCGCCCACGCGGTGCCGTTCTACGGGTGACCAATGAGAACTTTCCCTTCAATG	1436
47-6		1421
AV-1	ATTECCTCCACAGATCTCCATCTCCTCCCCCCCACACATCACCCCCAACCTCCT	1556
7A-3	.CT	1542
7A-6	•••••••••••••••••••••••••••••••••••••••	1541
	·	
AV-1	CCCCCCAGATCCACCCCCCCCCCCCCCCCCCCCCCCCCC	1676
YA - 5	.GAGTCAGAAA	1662
AV:0		
AV-1	TTCAACTCACCCGCCGTCTGGGGCATGACTTTGGCCAGGTGACAAGAGTCAAGAGTCAAGAGTTCTTCCGCTGGGGCGGAGGATCACGTGACGGAGGTGGCGGAGTGACTTCTACGTCAGAA	1796
AV-6	C.T. G. AA. GIT.	1781
-		
.v	P40/TATA P40/TATA **ACCOTOCIACIO COCCOCCOCCOCATCACCICCOCTATACACCICCOCCOCATCACTICCACCTCACCCCCACCTCACCCCCCACCCCCACCCCCCCC	1916
NV - 2		1009
V-6	G	1901

FIG 1B

AAV-	eccacasotaccaaaacaaatottectcotcacocogcatocttecagatoctotticcetocaagacatocagagaatgaatcagaatttcaccatttcattt	G 203
TYA-	l -A	201
YYA-(A	. 202
AAV-1	ACTOTTCMGAGTGCTTCCCCGGGGGTGTCAGAATCTCAACCGGTCGTCAGAAAGAGGACGTATCGGAAACTCTGTGCCCATTCATCATCTGCTGGGGGGGG	T 315
77A	T	
	A.TA.T	
	Rep/8 stop VP1 Rep68	stop
AAV-1	COSCUTIGORAL TOTAL CONTROL CON	G 227
AAV-6	AC	A 225: G 225:
 ∀-1	GGCATTCGCGAGTGGTGGGACTTGAAACCTGGAGCCCCGAAGCCCAAAGCCAAAGCAAGACAGCAAGAAG	
AAV-2	.A.AA.ACA.GC.CCC.A.ACCA.A.GC.GCAGGGC.TAA.A.TG.	2171
AAV-6	.A.AA.ACA.GC.CCC.A.ACCA.A.GC.GCAGGGC.TAA.ATG	. 2371
AAV-1	GGACTCGACAAGGGGGAGCCCOTCAACGGGGGACGCAGCGGGCCCTCGAGGACGACGACGACGAGGCCTACGACCAGCTCAAAGGGGGTGACAATCCGTACTGCGGTATAACCAGGCGAG	
AAV-2		
AAV-6		2476
AAV-1	GCCGAGTTTCAGGAGCGTCTGCAAGAAGATAGGTCTTTTGGGGGGCAACCTCGGGCGAGCAGTCTTCCAGGCCAAGAAGGGGGCTTCTGGAACCTCTGGGTTGAGGAAGGGGCTAAG	
AAV-6		2618
	VP2 ACCOUNTESCALAGALACTORICOGETAGAGCAGACTOCCACALGACCAGACTCCTCCCCCCACACACACCAGCAGCCCCCCTALALAGAGACTCAATTTTCGTCAGACTGGC	
AAV-6	TGAC.T	2738
117-1	CLUTTER CONTROL PROCESS AND CONTROL CASE A CONTROL	
MV-2	GACTCAGAGTCAGTCCCCGATCCACAACCTCTCGGAGAACCTCCAGCAACCCCCCGCTGTGTGGGACCTACTACGATTGGCTTCAGGCGTGGGGACCAATGGCAAGAACAATAACGAAGGCGGA.T.G.G.G.A.A.A.G.G.G.A.A.A.G.G.G.G	2873
MA-8	TGCCCAAG.ATAG	2858
LAV-1	GCCCACCCAGTAGCCTAATGCCTCAGGAAATTGGCATTGCGATTCCACATGGCTGGGGGACAGGTCATCACCACCAGGCACCTGGGCCCTGCGCACCTACAATAACCACCTCTAC	7001
LAV-2		2973
rv∆-4	T C	2978
AV-1	ALGCALATCTCCAGTGCTTCAACGGGGGCCAGCAACGACCACTACTTCGGCTACAGCACCCCCTGGGGGTATTTGATTTCAACAGATTCCACTGCCACTTTTCACCACGTGACTGG	3113
WA-3	ATCCAAAATCUT	
AV-1		3078
AV-1	CAGGGACTCATCAACAACTTGGGGGATTCCGGCCCCAAGAGACTCAACTTCAACTCTTCAACATCCAAGTCAAGGAGGTCACGACGAATGATGGCGTCACAACCATCGCTAATAACCTT	3233
AV-1	G	3218
AV-1	ACCASCACOGTICAAOTCTTCTCGGACTCGGAGTACCAGCTTCCGTACGTCCTCGGCACCAGCGCTCCCTCC	****
AV-6	T.G	3338
AV-1	TACCTCACCCTCAACAATGGCAGCCGAGCCGTTCATCCTTTTACTGCCTGGAATATTTCCCTTCTCAGATCCTCAGAACGGGCAACAACTTTACCTTCAGCTACACCTTTTAG	3671
AV - 2		
AV-6		3454
LV-1	CALGITOCCTTTCCACAGCAGCTACGCGCACAGCCAGAGCCTGGACGGCCTGATGAATCCTCTCATCGACCAATACCTGTATTACCTGAACAGAACTCAAAATCAGTCCGGAAGTGCCCAA	3597
LV - Z		
.V-6		3378
W-1	AACAAGGACTTGCTGTTTAGGCGTGGGTGTGCGATGTGTGTTCAGGCCAAAACTGGCTACCTGGACCCTGTTATCGGCAGGAGGGGGGTTTCTAAACAAAAACAGACAACAAC	3713
W-3	C.GTCAAGCC.T.ATCT.AG.CCCGAG.GAGATCGG.ACT.T.GG	
.v-€		3698
V-1	AACAGCAATTITACCTGGACTGGTGGTTCAAAATATAACCTCAATGGGGGTGAATCCATCATCAGCCTGGCACTGCTATGGCCTCACACAAGAGGACGAAGACGAAGTTCTTTCCCATG	1433
V-3	TG.A.ACT.GAA.CGCC	3010

FIG IC

444-1	ACCOSTETCATEATTTTTGGAAAAGAGGCCCCGGAGCTTCAAACACTGCATTGGACAATGATTACAGACCAAGAGGAAATTAAAGCCACTAAGCCCTGTGGCCAAGGATTT 1951
77A-9	GTC.CCGGC.AGT.A.AGAAAATGTGAACA.TAG
AAV-1	GGGACCGTGGCAGTCAATTTCCAGAGCAGCACAGACCCTGCGACCGGAGATGTGCATGCTATGGGAGCATTACCTGGCATGGTATGGCAAGATAGAGACGTGTACCTGCAGGGTCCC 4873
YYA - 3	TT.T.:AT.TACCCAGAG.C.AG.ATCC.CCC.
AAV-6	T
AAV-1	ATTTGGGCCAAAATTCCTCACACAGATGGACACTTTCACCCGTCTCCTCTTATGGGCGGCTTTTGGACTCAAGAACCCGCCTCCTCAGATCCTCATCAAAAAACACGCCTGTTCCTGCGAAT 4193
AAV-3	CAGAGCT
***-6	6178
AAV-1	CCTCCCCCCGGGGTTTTCACCTACAAGTTTCCTTCATTCA
AAV-3	TA.CACCCAGTGGC
AAV-6	AG
AAV-1	CGAAGTGCAGTACACATCCAATTATGCAAAATCTGCCAA-CGTTCATTTTACTGTGGACAACAATGGACTTTATACTGLGCCTCGCCCCATTGGCACCCGTTACCCTTACCCGTCCCCTGT 4431
MV-3	A.TTCCAACG.,TTTGCC. = ce con ce
M4-6	TTC
	VP1-3 stop PolyA signal
LAV-1	AATTACOTOTTAATCAATAAACCOGTTGAATCOTTTCAGTTGAACTTTGGTCTCCTGTCCTTATCTTAT
ware	1111Y-5-11
WAS-	07
AV-1	TTCGCGATAAAAGACTTACUTCATCGGGttacccctagtgatggagttgcccactccctctctgcgcgctcgct
74-6	######################################
AV-1	tggtccgcaggccccaccgagcgagcgagcgcagagagggagtgggcaa 4718
AV-2	ca.g.gctgt
AV-6	

aav-1 itr	AAV-1 ITR TCCATCACTAGGGG1
C C C C C C C C C C C C C C C C C C C	C GGCCCCACCGAGCGAGCGCGTTCTCTCCCTCACCGTTAAV-1 ITR C GC
A A G C G C G C G C G C G C G C G C G C	A CCGGGGTGGCTC C GCCCCACCGAG G C

AAV-1 TR






FIG. 3A










FIG. 4B



Group	1	2	3	4	5	6
Vector1 - α1AT						
Vector2-EPO	AAV2	AAV1	AAV2	AAV1	AAV1	AAV2



Group	1	2	3	4	5	6
Vector1 - α1AT	AAV2	AAV1	PBS	PBS	AAV2	AAV1
V ctor2-EPO	AAV2	AAV1	AAV2	AAV1	AAV1	AAV2

SEQUENCE LISTING

5

<110> Wilson, James M. Xiao, Weidong The Trustees of the University of Pennsylvania <120> Adeno-Associated Virus Serotype I Nucleic Acid Sequences, Vectors and Host Cells Containing Same <130> GNVPN.031PCT <140> <141> <150> 60/107,114 <151> 1998-11-05 <160> 20 <170> PatentIn Ver. 2.0 <210> 1 <211> 4718 <212> DNA <213> AAV-1 <220> <221> CDS <222> (335)..(2206) <220> <221> CDS <222> (2223)..(4430) <400> 1 ttgcccactc cctctctgcg cgctcgctcg ctcggtgggg cctgcggacc aaaggtccgc 60 agacggcaga gctctgctct gccggcccca ccgagcgagc gagcgcgcag agagggagtg 120 ggcaactcca tcactagggg taatcgcgaa gcgcctccca cgctgccgcg tcagcgctga 180 cgtaaattac gtcatagggg agtggtcctg tattagctgt cacgtgagtg cttttgcgac 240 attttgcgac accacgtggc catttagggt atatatggcc gagtgagcga gcaggatctc 300 cattttgacc gcgaaatttg aacgagcagc agcc atg ccg ggc ttc tac gag atc 355

1

Met Pro Gly Phe Tyr Glu Ile

	gtg Val								403
	gtg Val								451
	atg Met								499
	ctg Leu								547
	gag Glu 75								595
	ctc Leu					_		_	643
	cgc Arg								691
	ggg Gly						 		739
	aat Asn								787
	aac Asn 155								835
	aac Asn								883
	cgg Arg								931

									cct Pro	gtc Val 215	979
									 tgg Trp 230	_	1027
-				 -	-			_	 gac Asp		1075
							_		cag Gln		1123
								_	aaa Lys		1171
									aaa Lys		1219
									gcc Ala 310		1267
									aag Lys		1315
						-		_	aac Asn		1363
									aac Asn		1411
									gtg Val		1459
									gcc Ala 390		1507

gcc Ala	att	cto Leu	ggc Gly 395	Gly	ago Ser	aag Lys	gtg Val	Arg 400	Val	gac Asp	caa Glr	aag Lys	tgc Cys 405	Lys	tcg Ser	1555
tcc	gcc	Gln 410	Ile	gac Asp	CCC	acc Thr	Pro	Val	atc Ile	gto Val	acc Thr	Ser 420	Asn	acc Thr	aac Asn	1603
atg Met	tgc Cys 425	Ala	gtg Val	att Ile	gac Asp	ggg Gly 430	Asn	agc Ser	acc Thr	acc	tto Phe	Glu	Cac	cag Gln	cag Gln	1651
ccg Pro 440	ttg Leu	cag Gln	gac Asp	cgg Arg	atg Met 445	ttc Phe	aaa Lys	ttt Phe	gaa Glu	ctc Leu 450	Thr	cgc Arg	cgt Arg	ctg Leu	gag Glu 455	1699
cat His	gac Asp	ttt Phe	ggc	aag Lys 460	gtg Val	aca Thr	aag Lys	cag Gln	gaa Glu 465	gtc Val	aaa Lys	gag Glu	ttc Phe	ttc Phe 470	cgc Arg	1747
tgg Trp	gcg Ala	cag Gln	gat Asp 475	cac His	gtg Val	acc Thr	gag Glu	gtg Val 480	gcg Ala	cat His	gag Glu	ttc Phe	tac Tyr 485	gtc Val	aga Arg	1795
aag Lys	ggt Gly	gga Gly 490	gcc Ala	aac Asn	aaa Lys	aga Arg	ccc Pro 495	gcc Ala	ccc Pro	gat Asp	gac Asp	gcg Ala 500	gat Asp	aaa Lys	agc Ser	1843
gag Glu	ccc Pro 505	aag Lys	cgg Arg	gcc Ala	tgc Cys	ccc Pro 510	tca Ser	gtc Val	gcg Ala	gat Asp	cca Pro 515	tcg Ser	acg Thr	tca Ser	gac Asp	1891
			gct Ala													1939
tct Ser	cgt Arg	cac His	gcg Ala	ggc Gly 540	atg Met	ctt Leu	cag Gln	atg Met	ctg Leu 545	ttt Phe	ccc Pro	tgc Cys	aag Lys	aca Thr 550	tgc Cys	1987
gag Glu	aga Arg	atg Met	aat Asn 555	cag Gln	aat Asn	ttc Phe	aac Asn	att Ile 560	tgc Cys	ttc Phe	acg Thr	cac His	ggg Gly 565	acg Thr	aga Arg	2035
gac Asp	tgt Cys	tca Ser 570	gag Glu	tgc Cys	ttc Phe	Pro	ggc Gly 575	gtg Val	tca Ser	gaa Glu	tct Ser	caa Gln 580	ccg Pro	gtc Val	gtc Val	2083

_	-						-	•				-	ctg Leu	,,,,	2131
	-	_		-	-	-	-	-	-	-	-		gtg Val	_	2179
			gtt Val 620				taa	atg	actt	aaa (ccag	M		ct gcc la Ala	2231
_				_				-					ggc Gly		2279
													gcc Ala		2327
													tac Tyr	_	2375
												_	aac Asn 690		2423
											_	_	cag Gln		2471
													gcc Ala		2519
													ctc Leu		2567
													ggt Gly		2615
													gta Val 770		2663

						ggc	-		ggc Gly	2711
						act Thr 800		-		2759
						cca Pro	-			2807
						ggc Gly				2855
						gcc Ala				2903
						atc Ile			-	2951
						ctc Leu 880				2999
						cac His				3047
						ttc Phe				3095
						aat Asn				3143
						caa Gln				3191
						aac Asn 960				3239

acg gtt caa gtc ttc tcg gac tcg gag tac cag ctt ccg tac gtc ctc Thr Val Gln Val Phe Ser Asp Ser Glu Tyr Gln Leu Pro Tyr Val Leu 965 970 975	3287
ggc tct gcg cac cag ggc tgc ctc cct ccg ttc ccg gcg gac gtg ttc Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro Ala Asp Val Phe 980 985 990 995	3335
atg att ccg caa tac ggc tac ctg acg ctc aac aat ggc agc caa gcc Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn Gly Ser Gln Ala 1000 1005 1010	3383
gtg gga cgt toa too ttt tac tgc ctg gaa tat ttc cct tot cag atg Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe Pro Ser Gln Met 1015 1020 1025	3431
ctg aga acg ggc aac aac ttt acc ttc agc tac acc ttt gag gaa gtg Leu Arg Thr Gly Asn Asn Phe Thr Phe Ser Tyr Thr Phe Glu Glu Val 1030 1035 1040	3479
cct ttc cac age age tae geg cac age cag age ctg gac egg etg atg Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp Arg Leu Met 1045 1050 1055	3527
Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Asn Arg Thr Gln Asn 1060 1065 1070 1075	3575
Gln Ser Gly Ser Ala Gln Asn Lys Asp Leu Leu Phe Ser Arg Gly Ser 1080 1085 1090	3623
Pro Ala Gly Met Ser Val Gln Pro Lys Asn Trp Leu Pro Gly Pro Cys 1095 1100 1105	3671
Tyr Arg Gln Gln Arg Val Ser Lys Thr Lys Thr Asp Asn Asn Ser 1110 1115 1120	3719
Asn Phe Thr Trp Thr Gly Ala Ser Lys Tyr Asn Leu Asn Gly Arg Glu 1125 1130 1135	3767
tcc atc atc aac cct ggc act gct atg gcc tca cac aaa gac gac gaa Ser Ile Ile Asn Pro Gly Thr Ala Met Ala Ser His Lys Asp Asp Glu 1140 1145 1150 1155	3815

-	•		ttt Phe		-	-		Val	-				Lys		-	3863
-		Ala	tca Ser 175			-	Leu	-		-	-	Ile		-	-	3911
	Glu		aaa Lys	-		Asn			-		Glu	•				3959
Val			aat Asn		Gln					Asp					-	4007
	His		atg Met	Gly					Met	-			-	Arg	_	4055
			cag Gln 1					Ala					Thr	-		4103
		His	ccg Pro .255				Met					Leu	_		-	4151
	Pro		atc Ile			Lys ·					Pro	_			_	4199
Ala			tca Ser		Thr					Phe						4247
	Gly		gtg Val	Ser					Trp					Glu		4295
			tgg Trp 1					Gln					Tyr	-		4343
		Asn	gtt Val 335				Val					Leu				4391

cct cgc ccc att ggc acc cgt tac ctt acc cgt ccc ctg taattacgtg

Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu

1350

1355

1360

ttaatcaata aaccggttga ttcgtttcag ttgaactttg gtctcctgtc cttcttatct 4500
tatcggttac catggttata gcttacacat taactgcttg gttgcgcttc gcgataaaag 4560
acttacgtca tcgggttacc cctagtgatg gagttgccca ctccctctt gcgcgctcgc 4620
tcgctcggtg gggcctgcgg accaaaggtc cgcagacggc agagctctgc tctgccggcc 4680
ccaccgagcg agcgagcgcg cagagaggga gtgggcaa 4718

<210> 2

<211> 623

<212> PRT

<213> AAV-1

<400> 2

Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp 1 5 10 15

Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu 20 25 30

Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile 35 40 45

Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu 50 55 60

Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val 65 70 75 80

Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu 85 90 95

Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile 100 105 110

Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu 115 120 125

Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly 130 135 140

)"

Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys Thr Gln Pro Glu Leu Gln Trp Ala Trp Thr Asn Met Glu Glu Tyr Ile Ser Ala Cys Leu Asn Leu Ala Glu Arg Lys Arg Leu Val Ala Gln His Leu Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu Asn Leu Asn Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser Ala Arg Tyr Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu Asn Gly Tyr Glu Pro Ala Tyr Ala Gly Ser Val Phe Leu Gly Trp Ala ·Gln Lys Arg Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg

Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val 405 410 415

Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser 420 425 430

Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe 435 440 445

Glu Leu Thr Arg Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln 450 455 460

Glu Val Lys Glu Phe Phe Arg Trp Ala Gln Asp His Val Thr Glu Val 465 470 475 480

Ala His Glu Phe Tyr Val Arg Lys Gly Gly Ala Asn Lys Arg Pro Ala 485 490 495

Pro Asp Asp Ala Asp Lys Ser Glu Pro Lys Arg Ala Cys Pro Ser Val 500 505 510

Ala Asp Pro Ser Thr Ser Asp Ala Glu Gly Ala Pro Val Asp Phe Ala 515 520 525

Asp Arg Tyr Gln Asn Lys Cys Ser Arg His Ala Gly Met Leu Gln Met 530 540

Leu Phe Pro Cys Lys Thr Cys Glu Arg Met Asn Gln Asn Phe Asn Ile 545 550 555 560

Cys Phe Thr His Gly Thr Arg Asp Cys Ser Glu Cys Phe Pro Gly Val 565 570 575

Ser Glu Ser Gln Pro Val Val Arg Lys Arg Thr Tyr Arg Lys Leu Cys 580 585 590

Ala Ile His His Leu Leu Gly Arg Ala Pro Glu Ile Ala Cys Ser Ala 595 600 605

Cys Asp Leu Val Asn Val Asp Leu Asp Asp Cys Val Ser Glu Gln 610 615 620

<210> 3

<211> 736

<212> PRT

<213> AAV-1

<400>	3
-------	---

- Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser

 1 10 15
- Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro 20 25 30
- Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro 35 40 45
- Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro 50 55 60
- Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
 65 70 75 80
- Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala 85 90 95
- Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly
 100 105 110
- Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro 115 120 125
- Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Lys Arg 130 135 140
- Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ser Gly Ile Gly 145 150 155 160
- Lys Thr Gly Gln Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr 165 170 175
- Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro Pro 180 185 190
- Ala Thr Pro Ala Ala Val Gly Pro Thr Thr Met Ala Ser Gly Gly Gly 195 200 . 205
- Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn Ala 210 215 220
- Ser Gly Asn Trp His Cys Asp Ser Thr Trp Leu Gly Asp Arg Val Ile 225 230 235 240
- Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu

245 250 255

Tyr Lys Gln Ile Ser Ser Ala Ser Thr Gly Ala Ser Asn Asp Asn His 260 265 270

Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg Phe 275 280 285

His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn Asn 290 295 300

Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile Gln 305 310 315

Val Lys Glu Val Thr Thr Asn Asp Gly Val Thr Thr Ile Ala Asn Asn 325 330 335

Leu Thr Ser Thr Val Gln Val Phe Ser Asp Ser Glu Tyr Gln Leu Pro 340 345 350

Tyr Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro Ala 355 360 365

Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn Gly 370 375 380

Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe Pro 385 390 395 400

Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Thr Phe Ser Tyr Thr Phe 405 410 415

Glu Glu Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp 420 425 430

Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Asn Arg 435 440 445

Thr Gln Asn Gln Ser Gly Ser Ala Gln Asn Lys Asp Leu Leu Phe Ser 450 455 460

Arg Gly Ser Pro Ala Gly Met Ser Val Gln Pro Lys Asn Trp Leu Pro 465 470 475 480

Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Lys Thr Lys Thr Asp Asn 485 490 495

Asn Asn Ser Asn Phe Thr Trp Thr Gly Ala Ser Lys Tyr Asn Leu Asn

510

500 505

Gly Arg Glu Ser Ile Ile Asn Pro Gly Thr Ala Met Ala Ser His Lys 515 520 525

Asp Asp Glu Asp Lys Phe Phe Pro Met Ser Gly Val Met Ile Phe Gly 530 535 540

Lys Glu Ser Ala Gly Ala Ser Asn Thr Ala Leu Asp Asn Val Met Ile 545 550 555 560

Thr Asp Glu Glu Ile Lys Ala Thr Asn Pro Val Ala Thr Glu Arg 565 570 575

Phe Gly Thr Val Ala Val Asn Phe Gln Ser Ser Ser Thr Asp Pro Ala 580 585 590

Thr Gly Asp Val His Ala Met Gly Ala Leu Pro Gly Met Val Trp Gln 595 600 605

Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His 610 615 620

Thr Asp Gly His Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Leu 625 630 635 640

Lys Asn Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala 645 650 655

Asn Pro Pro Ala Glu Phe Ser Ala Thr Lys Phe Ala Ser Phe Ile Thr 660 665 670

Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln 675 680 685

Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Val Gln Tyr Thr Ser Asn 690 695 700

Tyr Ala Lys Ser Ala Asn Val Asp Phe Thr Val Asp Asn Asn Gly Leu 705 710 715 720

Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu
725 730 735

<210> 4

<211> 1872

<212> DNA

<213> AAV-1

<220> <221> CDS <222> (1)..(1869)

<400> 4

atg ccg ggc ttc tac gag atc gtg atc aag gtg ccg agc gac ctg gac

Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp

1 5 10 15

gag cac ctg ccg ggc att tct gac tcg ttt gtg agc tgg gtg gcc gag 96 Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu 20 .25 30

aag gaa tgg gag ctg ccc ccg gat tct gac atg gat ctg aat ctg att 144
Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile
35 40 45

gag cag gca ccc ctg acc gtg gcc gag aag ctg cag cgc gac ttc ctg 192 Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu 50 55 60

gtc caa tgg cgc cgc gtg agt aag gcc ccg gag gcc ctc ttc ttt gtt 240

Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val

65 70 75 80

cag ttc gag aag ggc gag tcc tac ttc cac ctc cat att ctg gtg gag 288
Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu
85 90 95

acc acg ggg gtc aaa tcc atg gtg ctg ggc cgc ttc ctg agt cag att 336
Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile
100 105 110

agg gac aag ctg gtg cag acc atc tac cgc ggg atc gag ccg acc ctg

Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu

115

120

125

ccc aac tgg ttc gcg gtg acc aag acg cgt aat ggc gcc gga ggg ggg 432 Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly Gly 130 135 140

aac aag gtg gtg gac gag tgc tac atc ccc aac tac ctc ctg ccc aag480Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys145150

act cag ccc gag ctg cag tgg gcg tgg act aac atg gag gag tat ata 528

	•																
	Thr	Gln	Pro	Glu	Leu 165	Gln	Trp	Ala	Trp	Thr 170	Asn	Met	Glu	Glu	Туг 175	Ile	
															cag Gln		576
															ctg Leu		624
															cgc Arg		672
															gag Glu		720
															gcc Ala 255		768
															ggc Gly		816
															ccc Pro		864
															gag Glu		912
															tgg Trp		960
									Thr						ccg Pro 335		1008
								Ala							gtg Val		1056
1	ttc	tac	ggc	tgc	gtc	aac	tgg	acc	aat	gag	aac	ttt	ccc	ttc	aat	gat	1104

Phe	Туr	G1 y 355		Val	Asn	Trp	Thr 360		Glu	Asn	Phe	9ro		Asn	Asp	
					gtg Val							Lys				1152
					gcc Ala 390											1200
					aag Lys											1248
					acc Thr											1296
					cag Gln											1344
					ctg Leu											1,392
					ttc Phe 470											1440
					gtc Val											1488
					aaa Lys											1536
gcg Ala	gat Asp	cca Pro 515	tcg Ser	acg Thr	tca Ser	gac Asp	gcg Ala 520	gaa Glu	gga Gly	gct Ala	ccg Pro	gtg Val 525	gac Asp	ttt Phe	gcc Ala	1584
					aaa Lys											1632
ctg	ttt	ccc	tgc	aag	aca	tgc	gag	aga	atg	aat	cag	aat	ttc	aac	att	1680

Leu Phe Pro Cys Lys Thr Cys Glu Arg Met Asn Gln Asn Phe Asn Ile 545 550 555 560

tgc ttc acg cac ggg acg aga gac tgt tca gag tgc ttc ccc ggc gtg 1728 Cys Phe Thr His Gly Thr Arg Asp Cys Ser Glu Cys Phe Pro Gly Val 565 570 575

tca gaa tct caa ccg gtc gtc aga aag agg acg tat cgg aaa ctc tgt 1776 Ser Glu Ser Gln Pro Val Val Arg Lys Arg Thr Tyr Arg Lys Leu Cys 580 585 590

gcc att cat ctg ctg ggg cgg gct ccc gag att gct tgc tcg gcc 1824 Ala Ile His His Leu Leu Gly Arg Ala Pro Glu Ile Ala Cys Ser Ala 595 600 605

tgc gat ctg gtc aac gtg gac ctg gat gac tgt gtt tct gag caa taa 1872 Cys Asp Leu Val Asn Val Asp Leu Asp Asp Cys Val Ser Glu Gln 610 615 620

<210> 5 <211> 623 <212> PRT <213> AAV-1

<400> 5

Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp

1 10 15

Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu 20 25 30

Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile $35 \hspace{1cm} 40 \hspace{1cm} 45$

Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu 50 55 60

Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val 65 70 75 80

Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu 85 90 95

Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile 100 .105 110

Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu $\,\cdot\,$

Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys Thr Gln Pro Glu Leu Gln Trp Ala Trp Thr Asn Met Glu Glu Tyr Ile Ser Ala Cys Leu Asn Leu Ala Glu Arg Lys Arg Leu Val Ala Gln His Leu Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu Asn Leu Asn Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser Ala Arg Tyr Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu Asn Gly Tyr Glu Pro Ala Tyr Ala Gly Ser Val Phe Leu Gly Trp Ala Gln Lys Arg Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala

Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro 340 345 350

Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp 355 360 365

Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala

370 375 380

Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg 385 390 395 400

- Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val 405 410 415
- Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser
 420 425 430
- Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe 435 440 445
- Glu Leu Thr Arg Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln 450 455 460
- Glu Val Lys Glu Phe Phe Arg Trp Ala Gln Asp His Val Thr Glu Val 465 470 475 480
- Ala His Glu Phe Tyr Val Arg Lys Gly Gly Ala Asn Lys Arg Pro Ala 485 490 495
- Pro Asp Asp Ala Asp Lys Ser Glu Pro Lys Arg Ala Cys Pro Ser Val 500 505 510
- Ala Asp Pro Ser Thr Ser Asp Ala Glu Gly Ala Pro Val Asp Phe Ala 515 520 525
- Asp Arg Tyr Gln Asn Lys Cys Ser Arg His Ala Gly Met Leu Gln Met 530 540
- Leu Phe Pro Cys Lys Thr Cys Glu Arg Met Asn Gln Asn Phe Asn Ile 545 550 555 560
- Cys Phe Thr His Gly Thr Arg Asp Cys Ser Glu Cys Phe Pro Gly Val 565 570 575
- Ser Glu Ser Gln Pro Val Val Arg Lys Arg Thr Tyr Arg Lys Leu Cys 580 585 590
- Ala Ile His His Leu Leu Gly Arg Ala Pro Glu Ile Ala Cys Ser Ala 595 600 605
- Cys Asp Leu Val Asn Val Asp Leu Asp Asp Cys Val Ser Glu Gln 610 615 . 620

<210> 6 <211> 1641 <212> DNA <213> AAV-1 <220> <221> CDS <222> (1)..(1638) <400> 6 atg ccg ggc ttc tac gag atc gtg atc aag gtg ccg agc gac ctg gac 48 Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp 10 gag cac ctg ccg ggc att tct gac tcg ttt gtg agc tgg gtg gcc gag 96 Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu aag gaa tgg gag ctg ccc ccg gat tct gac atg gat ctg aat ctg att 144 Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile gag cag gca ccc ctg acc gtg gcc gag aag ctg cag cgc gac ttc ctg 192 Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu 50 55 gtc caa tgg cgc cgc gtg agt aag gcc ccg gag gcc ctc ttc ttt gtt 240 Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val 65 70 cag ttc gag aag ggc gag tcc tac ttc cac ctc cat att ctg gtg gag 288 Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu 85 90 acc acg ggg gtc aaa tcc atg gtg ctg ggc cgc ttc ctg agt cag att 336 Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile 100 110 agg gac aag ctg gtg cag acc atc tac cgc ggg atc gag ccg acc ctg 384 Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu 115 120 ccc aac tgg ttc gcg gtg acc aag acg cgt aat ggc gcc gga ggg ggg 432 Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly 130 135

21

aac aag gtg gtg gac gag tgc tac atc ccc aac tac ctc ctg ccc aag

Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys

480

150 155 160 act cag ccc gag ctg cag tgg gcg tgg act aac atg gag gag tat ata. Thr Gln Pro Glu Leu Gln Trp Ala Trp Thr Asn Met Glu Glu Tyr Ile 165 170 age gee tgt ttg aac etg gee gag ege aaa egg ete gtg geg eag eac Ser Ala Cys Leu Asn Leu Ala Glu Arg Lys Arg Leu Val Ala Gln His 180 ctg acc cac gtc agc cag acc cag gag cag aac aag gag aat ctg aac Leu Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu Asn Leu Asn 195 200 ccc aat tct gac gcg cct gtc atc cgg tca aaa acc tcc gcg cgc tac 672 Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser Ala Arg Tyr 210 215 atg gag ctg gtc ggg tgg ctg gtg gac cgg ggc atc acc tcc gag aag 720 Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys 225 230 235 cag tgg atc cag gag gac cag gcc tcg tac atc tcc ttc aac gcc gct 768 Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala 245 250 255 tcc aac tcg cgg tcc cag atc aag gcc gct ctg gac aat gcc ggc aag Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys 260 265 atc atg gcg ctg acc aaa tcc gcg ccc gac tac ctg gta ggc ccc gct 864 Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala 275 280 ccg ccc gcg gac att aaa acc aac cgc atc tac cgc atc ctg gag ctg 912 Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu 290 295 300 aac ggc tac gaa cct gcc tac gcc ggc tcc gtc ttt ctc ggc tgg gcc 960 Asn Gly Tyr Glu Pro Ala Tyr Ala Gly Ser Val Phe Leu Gly Trp Ala 305 310 320 cag aaa agg ttc ggg aag cgc aac acc atc tgg ctg ttt ggg ccg gcc 1008 Gln Lys Arg Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala 325 330

acc acg ggc aag acc aac atc gcg gaa gcc atc gcc cac gcc gtg ccc Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro

340 345 350

	c tac e Tyr							Asn					Phe			1104
	gtc Val 370	Asp										Lys				1152
	g gtc s Val															1200
	g gac . Asp															1248
	gtc Val															1296
acc Thi	acc Thr	ttc Phe 435	gag Glu	cac His	cag Gln	cag Gln	ccg Pro 440	ttg Leu	cag Gln	gac Asp	cgg Arg	atg Met 445	ttc Phe	aaa Lys	ttt Phe	1344
	ctc Leu 450															1392
	gtc Val															1440
gcg Ala	cat His	gag Glu	ttc Phe	tac Tyr 485	gtc Val	aga Arg	aag Lys	ggt Gly	gga Gly 490	gcc Ala	aac Asn	aaa Lys	aga Arg	ccc Pro 495	gcc Ala	1488
ccc Pro	gat Asp	gac Asp	gcg Ala 500	gat Asp	aaa Lys	agc Ser	gag Glu	ccc Pro 505	aag Lys	cgg Arg	gcc Ala	tgc Cys	ccc Pro 510	tca Ser	gtc Val	1536
	gat Asp															1584
gac Asp	agg Arg	tat Tyr	ggc Gly	tgc Cys	cga Arg	tgg Trp	tta Leu	tct Ser	tcc Ser	aga Arg	ttg Leu	gct	cga Arg	gga Gly	caa Gln	1632

530 535 540

cct ctc tga 1641 .

Pro Leu 545

<210> 7

<211> 546

<212> PRT

<213> AAV-1

<400> 7

Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp 1 5 10 15

Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu 20 25 30

Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile 35 40 45

Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu
50 55 60

Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val 65 70 75 80

Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu 85 90 95

Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile 100 105 110

Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu 115 120 125

Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly Gly 130 135 140

Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys 145 150 155 160

Thr Gln Pro Glu Leu Gln Trp Ala Trp Thr Asn Met Glu Glu Tyr Ile 165 170 175

Ser Ala Cys Leu Asn Leu Ala Glu Arg Lys Arg Leu Val Ala Gln His 180 185 190

Leu Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu Asn Leu Asn 195 200 205

- Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser Ala Arg Tyr 210 215 220
- Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys 225 230 235 240
- Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala 245 250 255
- Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys 260 265 270
- Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala 275 280 285
- Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu 290 295 300
- Asn Gly Tyr Glu Pro Ala Tyr Ala Gly Ser Val Phe Leu Gly Trp Ala 305 310 315 320
- Gln Lys Arg Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala 325 330 335
- Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro 340 345 350
- Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp 355 360 365
- Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala 370 375 380
- Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg 385 390 395 400
- Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val 405 410 415
- Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser 420 425 430
- Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe 435 440 445

Glu Leu Thr Arg Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln 455 Glu Val Lys Glu Phe Phe Arg Trp Ala Gln Asp His Val Thr Glu Val 470 475 Ala His Glu Phe Tyr Val Arg Lys Gly Gly Ala Asn Lys Arg Pro Ala 490 Pro Asp Asp Ala Asp Lys Ser Glu Pro Lys Arg Ala Cys Pro Ser Val 500 505 Ala Asp Pro Ser Thr Ser Asp Ala Glu Gly Ala Pro Val Asp Phe Ala 515 520 Asp Arg Tyr Gly Cys Arg Trp Leu Ser Ser Arg Leu Ala Arg Gly Gln 535 540 Pro Leu 545 <210> 8 <211> 1200 <212> DNA <213> AAV-1 <220> <221> CDS <222> (1)..(1197) <400> 8 atg gag ctg gtc ggg tgg ctg gtg gac cgg ggc atc acc tcc gag aag Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys 1 5 cag tgg atc cag gag gac cag gcc tcg tac atc tcc ttc aac gcc gct 96 Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala 20 25 tcc aac tcg cgg tcc cag atc aag gcc gct ctg gac aat gcc ggc aag 144 Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys 35 40 atc atg gcg ctg acc aaa tcc gcg ccc gac tac ctg gta ggc ccc gct Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala 50 . 55

			gac Asp													240
			gaa Glu													288
			ttc Phe 100													336
	_		aag Lys					-	-		_		_			384
			tgc Cys	-												432
_	-	-	aag Lys	_									-	_		480
_	-		gag Glu		-	_	-									528
	-		aag Lys 180	_	_	-						,				576
	-		tcc Ser				-	-	-			-			-	624
			gag Glu													672
-			cgc Arg													720
_	_		gag Glu			-			-	-						768

			gtc Val					-	816
			aaa Lys						864
			tca Ser						912
			aaa Lys 310						960
			aca Thr						1008
			acg Thr						1056
			gtc Val						1104
			ctg Leu						1152
			gtg Val 390					taa	1200
-210	\> 0								

<210> 9

<211> 399

<212> PRT

<213> AAV-1

<400> 9

Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys $1 \hspace{1cm} 5 \hspace{1cm} 10 \hspace{1cm} 15$

Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala

20

25

30

Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys 35 40 45

Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala 50 55 60

Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu 65 70 75 80

Asn Gly Tyr Glu Pro Ala Tyr Ala Gly Ser Val Phe Leu Gly Trp Ala 85 90 95

Gln Lys Arg Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala 100 105 110

Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro 115 120 125

Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp 130 135 140

Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala 145 150 155 160

Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg 165 170 175

Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val 180 185 190

Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser 195 200 205

Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe 210 215 220

Glu Leu Thr Arg Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln 225 230 235 240

Glu Val Lys Glu Phe Phe Arg Trp Ala Gln Asp His Val Thr Glu Val 245 250 255

Ala His Glu Phe Tyr Val Arg Lys Gly Gly Ala Asn Lys Arg Pro Ala 260 265 270

Pro Asp Asp Ala Asp Lys Ser Glu Pro Lys Arg Ala Cys Pro Ser Val

275 280 285

Ala Asp Pro Ser Thr Ser Asp Ala Glu Gly Ala Pro Val Asp Phe Ala 290 295 300

Asp Arg Tyr Gln Asn Lys Cys Ser Arg His Ala Gly Met Leu Gln Met 305 310 315 320

Leu Phe Pro Cys Lys Thr Cys Glu Arg Met Asn Gln Asn Phe Asn Ile
325 330 335

Cys Phe Thr His Gly Thr Arg Asp Cys Ser Glu Cys Phe Pro Gly Val 340 345 350

Ser Glu Ser Gln Pro Val Val Arg Lys Arg Thr Tyr Arg Lys Leu Cys 355 360 365

Ala Ile His His Leu Leu Gly Arg Ala Pro Glu Ile Ala Cys Ser Ala 370 380

Cys Asp Leu Val Asn Val Asp Leu Asp Asp Cys Val Ser Glu Gln 385 390 395

<210> 10

<211> 969

<212> DNA

<213> AAV-1

<220>

<221> CDS

<222> (1)..(966)

<220>

<221> misc_feature

<222> (943)..(944)

<223> minor splice site

<400> 10

atg gag ctg gtc ggg tgg ctg gtc ggc atc acc tcc gag aag 48
Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys
1 5 10 15

tcc aac tcg cgg tcc cag atc aag gcc gct ctg gac aat gcc ggc aag 144

Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys 35 40 atc atg gcg ctg acc aaa tcc gcg ccc gac tac ctg gta ggc ccc gct 192 Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala ccg ccc gcg gac att aaa acc aac cgc atc tac cgc atc ctg gag ctg 240 Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu 70 7:5 aac ggc tac gaa cct gcc tac gcc ggc tcc gtc ttt ctc ggc tgg gcc 288 Asn Gly Tyr Glu Pro Ala Tyr Ala Gly Ser Val Phe Leu Gly Trp Ala 85 cag aaa agg ttc ggg aag cgc aac acc atc tgg ctg ttt ggg ccg gcc 336 Gln Lys Arg Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala 100 105 acc acg ggc aag acc aac atc gcg gaa gcc atc gcc cac gcc gtg ccc Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro 115 120 ttc tac ggc tgc gtc aac tgg acc aat gag aac ttt ccc ttc aat gat 432 Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp 130 135 tgc gtc gac aag atg gtg atc tgg tgg gag gag ggc aag atg acg gcc 480 Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala 145 150 155 aag gtc gtg gag tcc gcc aag gcc att ctc ggc ggc agc aag gtg cgc 528 Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg 165 170 175 gtg gac caa aag tgc aag tcg tcc gcc cag atc gac ccc acc ccc gtg Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val 180 185 atc gtc acc tcc aac acc aac atg tgc gcc gtg att gac ggg aac agc 624 Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser 195 200 acc acc ttc gag cac cag cag ccg ttg cag gac cgg atg ttc aaa ttt 672 Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe 210 215 gaa ctc acc cgc cgt ctg gag cat gac ttt ggc aag gtg aca aag cag 720

WO 00/28061 PCT/US99/25694 Glu Leu Thr Arg Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln . 230 235 gaa gtc aaa gag ttc ttc cgc tgg gcg cag gat cac gtg acc gag gtg 768 Glu Val Lys Glu Phe Phe Arg Trp Ala Gln Asp His Val Thr Glu Val 245 250 gcg cat gag ttc tac gtc aga aag ggt gga gcc aac aaa aga ccc gcc 816 Ala His Glu Phe Tyr Val. Arg Lys Gly Gly Ala Asn Lys Arg Pro Ala 260 265 270 ccc gat gac gcg gat aaa agc gag ccc aag cgg gcc tgc ccc tca gtc 864 Pro Asp Asp Ala Asp Lys Ser Glu Pro Lys Arg Ala Cys Pro Ser Val 275 280 gcg gat cca tcg acg tca gac gcg gaa gga gct ccg gtg gac ttt gcc 912 Ala Asp Pro Ser Thr Ser Asp Ala Glu Gly Ala Pro Val Asp Phe Ala 290 295 gac agg tat ggc tgc cga tgg tta tct tcc aga ttg gct cga gga caa Asp Arg Tyr Gly Cys Arg Trp Leu Ser Ser Arg Leu Ala Arg Gly Gln 310 315 320 cct ctc tga 969 Pro Leu <210> 11 <211> 322 <212> PRT <213> AAV-1 <400> 11 Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala 25 Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys. 35 40 Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala 50

70

Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu

75

Asn	GIĀ	Tyr	GIU	85	АІа	Tyr	Ala	GIÀ	Ser 90		Phe	Leu	GIÀ	7rp 95	Ala
Gln	Lys	Arg	Phe 100	Gly	Lys	Arg	Asn	Thr 105		Trp	Leu	Phe	Gly 110	Pro	Ala
Thr	Thr	Gly 115	Lys	Thr	Asn	Ile	Ala 120	Glu	Ala	Ile	Ala	His 125		Val	Pro
Phe	Туг 130	Gly	Cys	Val	Asn	Trp 135	Thr	Asn	Glu	Asn	Phe 140		Phe	Asn	Asp
Cys 145	Val	Asp	Lys	Met	Val 150	Ile	Trp	Trp	Glu	Glu 155	Gly	Lys	Met	Thr	Ala 160
Lys	Val	Val	Glu	Ser 165	Ala	Lys	Ala	Ile	Leu 170	Gly	Gly	Ser	Lys	Val 175	Arg
Val	Asp	Gln	Lys 180	Cys	Lys	Ser	Ser	Ala 185	Gln	Ile	Asp	Pro	Thr 190	Pro	Val
Ile	Val	Thr 195	Ser	Asn	Thr	Asn	Met 200	Cys	Ala	Val	Ile	Asp 205	Gly	Asn	Ser
Thr	Thr 210	Phe	Glu	His	Gln	Gln 215	Pro	Leu	Gln	Asp	Arg 220	Met	Phe	Lys	Phe
G1u 225	Leu	Thr	Arg	Arg	Leu 230	Glu	His	Asp	Phe	Gly 235	Lys	Val	Thr	Lys	Gln 240
Glu	Val	Lys	Glu	Phe 245	Phe	Arg	Trp	Ala	Gln 250	Asp	His	Val	Thr	Glu 255	Val
Ala	His	Glu	Phe 260	Tyr	Val	Arg	Lys	Gly 265	Gly	Ala	Asn	Lys	Arg 270	Pro	Ala
Pro	Asp	Asp 275	Ala	Asp	Lys	Ser	Glu 280	Pro	Lys	Arg	Ala	Cys 285	Pro	Ser	Val
Ala	Asp 290	Pro	Ser	Thr	Ser	Asp 295	Ala	Glu	Gly	Ala	Pro 300	Val	Asp	Phe	Ala
Asp 305	Arg	Tyr	Gly	Cys	Arg 310	Trp	Leu	Ser	Ser	Arg 315	Leu	Ala	Arg	Gly	Gln 320
Pro	Leu														

<21 <21	0> 1 1> 2 2> D 3> A	211														
	1> C		(220	8)												
<40	0> 1	2														
					Tyr					Leu	gag Glu				Ser	48
														Lys	ccc Pro	96
								Asp			ggt Gly					144
											gac Asp 60					192
											gac Asp					240
											cgg Arg					288
gac Asp	gcc Ala	gag Glu	ttt Phe 100	cag Gln	gag Glu	cgt Arg	ctg Leu	caa Gln 105	gaa Glu	gat Asp	acg Thr	tct Ser	ttt Phe 110	el A aaa	ggc Gly	336
aac Asn	ctc Leu	ggg Gly 115	cga Arg	gca Ala	gtc Val	ttc Phe	cag Gln 120	gcc Ala	aag Lys	aag Lys	cgg Arg	gtt Val 125	ctc Leu	gaa Glu	cct Pro	384
ctc Leu	ggt Gly 130	ctg Leu	gtt Val	gag Glu	gaa Glu	ggc Gly 135 _.	gct Ala	aag Lys	acg Thr	gct Ala	cct Pro 140	gga Gly	aag Lys	aaa Lys	cgt Arg	432
ccg	gta	gag	cag	tcg	сса	caa	gag	cca	gac	tcc	tcc	tcg	qqc	atc	ggc	480

Pro 145	Val	Glu	Gln	Ser	Pro 150	Gln	Glu	Pro	Asp	Ser 155	Ser	Ser	Gly	Ile	Gly 160	
										Leu				cag Gln 175		528
									Gln					cct Pro		576
														ggt Gly		624
														aat Asn		672
														gtc Val		720
														cac His 255		768
														aac Asn		816
														aga Arg		864
cac His	tgc Cys 290	cac His	ttt Phe	tca Ser	cca Pro	cgt Arg 295	gac Asp	tgg Trp	cag Gln	cga Arg	ctc Leu 300	atc Ile	aac Asn	aac Asn	aat Asn	912
														atc Ile		960
														aat Asn 335		1008
ctt	acc	agc	acg	gtt	caa	gtc	ttc	tcg	gac	tcg	gag	tac	cag	ctt	ccg	1056

ţ,

Leu	Thr	Ser	Thr 340	Val	Gln	Val	Phe	ser 345	Asp	Ser	Glu	Tyr	Gln 350	Leu	Pro	
	_						_		-			-		ccg Pro		1104
-			-		_					-	_			aat Asn		1152
-		-			-					-	_	-		ttc Phe		1200
	-	-	-	-	-							-		acc Thr 415		1248
	-					-	_				_		-	ctg Leu	-	1296
	-	-					-			_			-	aac Asn	•	1344
			-		_						-			ttt Phe	-	1392
-				-		-		-	_					cta Leu		1440
		-			-	_	•	-						gac Asp 495		1488
														ctc Leu		1536
	_	_												cac His		1584
gac	gac	gaa	gac	aag	ttc	ttt	ccc	atg	agc	ggt	gtc	atg	att	ttt	gga	1632

Asp	Asp 530	Glu	Asp	Lys	Phe	Phe 535	Pro	Met	Ser	Gly	Val 540	Met	Ile	Phe	Gly	
		-	-	• •	gct Ala 550				-	-	-		-	-		1680
	-	_		-	att Ile		-					-		-	-	1728
				-	gtc Val			_	-	_	_		-		-	1776
					gct Ala							_				1824
					ctg Leu						-					1872
					cac His 630											1920
					cag Gln								-			1968
		_			ttt Phe		-		-		-					2016
					caa Gln		-		-		-		, ,	_	-	2064
					cgc Arg					_	_					2112
					aac Asn 710						-					2160
tat	act	gag	cct	cgc	ссс	att	ggc	acc	cgt	tac	ctt	acc	cgt	ccc	ctg	2208

Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu
725 730 735

taa 2211

<210> 13

<211> 736

<212> PRT

<213> AAV-1

<400> 13

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser

1 5 10 15

Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro 20 25 30

Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro 35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro 50 55 60

Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp 65 70 75 80

Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala 85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly
100 105 110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro 115 120 125

Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Lys Arg 130 135 140

Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Gly Ile Gly 145 150 155 160

Lys Thr Gly Gln Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr
165 170 175

Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro Pro 180 185 190 Ala Thr Pro Ala Ala Val Gly Pro Thr Thr Met Ala Ser Gly Gly Gly 195 200 205

3

- Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn Ala 210 215 220
- Ser Gly Asn Trp His Cys Asp Ser Thr Trp Leu Gly Asp Arg Val Ile 225 230 235 240
- Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu 245 250 255
- Tyr Lys Gln Ile Ser Ser Ala Ser Thr Gly Ala Ser Asn Asp Asn His 260 265 270
- Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg Phe 275 280 285
- His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn Asn 290 295 300
- Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile Gln 305 310 315 320
- Val Lys Glu Val Thr Thr Asn Asp Gly Val Thr Thr Ile Ala Asn Asn 325 330 335
- Leu Thr Ser Thr Val Gln Val Phe Ser Asp Ser Glu Tyr Gln Leu Pro 340 345 350
- Tyr Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro Ala 355 360 365
- Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn Gly 370 375 380
- Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe Pro 385 390 395 400
- Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Thr Phe Ser Tyr Thr Phe 405 410 415
- Glu Glu Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp 420 425 430
- Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Asn Arg
 435
 440
 445

Thr Gln Asn Gln Ser Gly Ser Ala Gln Asn Lys Asp Leu Leu Phe Ser 450 . 455 . 460

- Arg Gly Ser Pro Ala Gly Met Ser Val Gln Pro Lys Asn Trp Leu Pro 465 470 475 480
- Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Lys Thr Lys Thr Asp Asn 485 490 495
- Asn Asn Ser Asn Phe Thr Trp Thr Gly Ala Ser Lys Tyr Asn Leu Asn 500 505 510
- Gly Arg Glu Ser Ile Ile Asn Pro Gly Thr Ala Met Ala Ser His Lys 515 520 525
- Asp Asp Glu Asp Lys Phe Phe Pro Met Ser Gly Val Met Ile Phe Gly 530 540
- Lys Glu Ser Ala Gly Ala Ser Asn Thr Ala Leu Asp Asn Val Met Ile 545 550 555 560
- Thr Asp Glu Glu Ile Lys Ala Thr Asn Pro Val Ala Thr Glu Arg
 565 570 575
- Phe Gly Thr Val Ala Val Asn Phe Gln Ser Ser Ser Thr Asp Pro Ala 580 585 590
- Thr Gly Asp Val His Ala Met Gly Ala Leu Pro Gly Met Val Trp Gln 595 600 605
- Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His 610 615 620
- Thr Asp Gly His Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Leu 625 635 640
- Lys Asn Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala 645 650 655
- Asn Pro Pro Ala Glu Phe Ser Ala Thr Lys Phe Ala Ser Phe Ile Thr 660 665 670
- Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln 675 680 685
- Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Val Gln Tyr Thr Ser Asn 690 695 700

Tyr Ala Lys Ser Ala Asn Val Asp Phe Thr Val Asp Asn Asn Gly Leu 705 710 715 Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu 725 730 <210> 14 <211> 1800 <212> DNA <213> AAV-1 <220> <221> CDS <222> (1) .. (1797) <400> 14 acg gct cct gga aag aaa cgt ccg gta gag cag tcg cca caa gag cca Thr Ala Pro Gly Lys Lys Arg Pro Val Glu Gln Ser Pro Gln Glu Pro 10 gac tcc tcc tcg ggc atc ggc aag aca ggc cag cag ccc gct aaa aag Asp Ser Ser Ser Gly Ile Gly Lys Thr Gly Gln Gln Pro Ala Lys Lys. 20 aga ctc aat ttt ggt cag act ggc gac tca gag tca gtc ccc gat cca Arg Leu Asn Phe Gly Gln Thr Gly Asp Ser Glu Ser Val Pro Asp Pro 40 caa cct ctc gga gaa cct cca gca acc ccc gct gct gtg gga cct act 192 Gln Pro Leu Gly Glu Pro Pro Ala Thr Pro Ala Ala Val Gly Pro Thr 50 55 60 aca atg gct tca ggc ggt ggc gca cca atg gca gac aat aac gaa ggc Thr Met Ala Ser Gly Gly Gly Ala Pro Met Ala Asp Asn Asn Glu Gly 65 70 80 gcc gac gga gtg ggt aat gcc tca gga aat tgg cat tgc gat tcc aca Ala Asp Gly Val Gly Asn Ala Ser Gly Asn Trp His Cys Asp Ser Thr 85 90 tgg ctg ggc gac aga gtc atc acc acc agc acc cgc acc tgg gcc ttg Trp Leu Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu 100 105 110 ccc acc tac aat aac cac ctc tac aag caa atc tcc agt gct tca acg Pro Thr Tyr Asn Asn His Leu Tyr Lys Gln Ile Ser Ser Ala Ser Thr

120

115

		agc Ser														432
		gat Asp			-										tgg, Trp 160	480
	-	ctc Leu										-	-			528
		ctc Leu					-	•		-	•	-		-		576
_		acc Thr 195		-					_	_	_		_		-	624
-	-	gag Glu		-		-		-								672
-		cct Pro	_		•		•			Ξ.		-				720
	-	acg Thr					-		-			-				768
	-	ctg Leu	•					_	-	•	-	-				816
		ttc Phe 275	-										-			864
		agc Ser	-	-	-	-		-	_					-		912
	_	tat Tyr		-		-		-		_			-	-		960

_	-	-	ctg Leu 325	-	-				-		_		-	1008
			tgg Trp				_				_	-	-	1056
			aca Thr				_							1104
			aac Asn											1152
			tca Ser			_	_	_	_				-	1200
			att Ile 405											1248
			gtc Val						-			_		1296
			acc Thr										-	1344
			gac Asp							-	_		-	1392
			gtg Val											1440
			att Ile 485											1488
			ttt Phe							_				1536

aaa aac acg cct gtt cct gcg aat cct ccg gcg gag ttt tca gct aca Lys Asn Thr Pro Val Pro Ala Asn Pro Pro Ala Glu Phe Ser Ala Thr 515 520 aag ttt gct tca ttc atc acc caa tac tcc aca gga caa gtg agt gtg 1632 Lys Phe Ala Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Ser Val 535 540 gaa att gaa tgg gag ctg cag aaa gaa aac agc aag cgc tgg aat ccc Glu Ile Glu Trp Glu Leu Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro 545 550 555 gaa gtg cag tac aca tcc aat tat gca aaa tct gcc aac gtt gat ttt 1728 Glu Val Gln Tyr Thr Ser Asn Tyr Ala Lys Ser Ala Asn Val Asp Phe 565 570 act gtg gac aac aat gga ctt tat act gag cct cgc ccc att ggc acc Thr Val Asp Asn Asn Gly Leu Tyr Thr Glu Pro Arg Pro Ile Gly Thr 580 585 cgt tac ctt acc cgt ccc ctg taa 1800 Arg Tyr Leu Thr Arg Pro Leu 595 <210> 15 <211> 599 <212> PRT <213> AAV-1 <400> 15 Thr Ala Pro Gly Lys Lys Arg Pro Val Glu Gln Ser Pro Gln Glu Pro ⁻5 Asp Ser Ser Ser Gly Ile Gly Lys Thr Gly Gln Gln Pro Ala Lys Lys 20 Arg Leu Asn Phe Gly Gln Thr Gly Asp Ser Glu Ser Val Pro Asp Pro

Gln Pro Leu Gly Glu Pro Pro Ala Thr Pro Ala Ala Val Gly Pro Thr 50 55 60

Thr Met Ala Ser Gly Gly Gly Ala Pro Met Ala Asp Asn Asn Glu Gly 65 70 75 80

Ala Asp Gly Val Gly Asn Ala Ser Gly Asn Trp His Cys Asp Ser Thr

85 90 95

Trp Leu Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu 100 105 110

Pro Thr Tyr Asn Asn His Leu Tyr Lys Gln Ile Ser Ser Ala Ser Thr 115 120 125

Gly Ala Ser Asn Asp Asn His Tyr Phe Gly Tyr Ser Thr Pro Trp Gly
130 135 140

Tyr Phe Asp Phe Asn Arg Phe His Cys His Phe Ser Pro Arg Asp Trp 145 150 155 160

Gln Arg Leu Ile Asn Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn 165 170 175

Phe Lys Leu Phe Asn Ile Gln Val Lys Glu Val Thr Thr Asn Asp Gly
180 185 190

Val Thr Thr Ile Ala Asn Asn Leu Thr Ser Thr Val Gln Val Phe Ser 195 200 205

Asp Ser Glu Tyr Gln Leu Pro Tyr Val Leu Gly Ser Ala His Gln Gly 210 215 220

Cys Leu Pro Pro Phe Pro Ala Asp Val Phe Met Ile Pro Gln Tyr Gly 225 230 235 240

Tyr Leu Thr Leu Asn Asn Gly Ser Gln Ala Val Gly Arg Ser Ser Phe 245 250 255

Tyr Cys Leu Glu Tyr Phe Pro Ser Gln Met Leu Arg Thr Gly Asn Asn 260 265 270

Phe Thr Phe Ser Tyr Thr Phe Glu Glu Val Pro Phe His Ser Ser Tyr 275 280 285

Ala His Ser Gln Ser Leu Asp Arg Leu Met Asn Pro Leu Ile Asp Gln 290 295 300

Tyr Leu Tyr Tyr Leu Asn Arg Thr Gln Asn Gln Ser Gly Ser Ala Gln 305 310 315 320

Asn Lys Asp Leu Leu Phe Ser Arg Gly Ser Pro Ala Gly Met Ser Val 325 330 335

Gln Pro Lys Asn Trp Leu Pro Gly Pro Cys Tyr Arg Gln Gln Arg Val

340

345

350

Ser Lys Thr Lys Thr Asp Asn Asn Ser Asn Phe Thr Trp Thr Gly 355 360 365

Ala Ser Lys Tyr Asn Leu Asn Gly Arg Glu Ser Ile Ile Asn Pro Gly 370 375 380

Thr Ala Met Ala Ser His Lys Asp Asp Glu Asp Lys Phe Phe Pro Met 385 390 395 400

Ser Gly Val Met Ile Phe Gly Lys Glu Ser Ala Gly Ala Ser Asn Thr 405 410 415

Ala Leu Asp Asn Val Met Ile Thr Asp Glu Glu Glu Ile Lys Ala Thr 420 425 430

Asn Pro Val Ala Thr Glu Arg Phe Gly Thr Val Ala Val Asn Phe Gln 435 ' 440 445

Ser Ser Ser Thr Asp Pro Ala Thr Gly Asp Val His Ala Met Gly Ala 450 455 460

Leu Pro Gly Met Val Trp Gln Asp Arg Asp Val Tyr Leu Gln Gly Pro 465 470 475 480

Ile Trp Ala Lys Ile Pro His Thr Asp Gly His Phe His Pro Ser Pro
485 490 495

Leu Met Gly Gly Phe Gly Leu Lys Asn Pro Pro Pro Gln Ile Leu Ile 500 505 510

Lys Asn Thr Pro Val Pro Ala Asn Pro Pro Ala Glu Phe Ser Ala Thr 515 520 525

Lys Phe Ala Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Ser Val 530 535 540

Glu Ile Glu Trp Glu Leu Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro 545 550 555 560

Glu Val Gln Tyr Thr Ser Asn Tyr Ala Lys Ser Ala Asn Val Asp Phe
565 570 575

Thr Val Asp Asn Gly Leu Tyr Thr Glu Pro Arg Pro Ile Gly Thr 580 585 590

Arg Tyr Leu Thr Arg Pro Leu

595

	1> 1															
	1> 1															
	2> D															
\ 21	3> A	AV-1	•													
<22	2.															
	1> C	יחפ														
			(160	121												
122	٠, ۲	+/	1100	, ,												
<40	0> 1	6														
			ggc	aat	aac	αca	CCa	ato		420	22+	226	~		gcc	40
															Ala	48
1			1	5		n_u	110	ricc	10	лэр	ASII	ASII	GIU	_		
Ī				Ŭ					10					15		
gac	gga	ata	aat	aat	acc	tca	aaa	aat	taa	cat	tac	ast	taa	202	tgg	0.6
															Trp	96
			20				OL y	25	1-1	1113	Cys	ASP	30	1111	rrp	
								23					30			
ctg	qqc	gac	aga	atc	atc	acc	acc	agc	acc	cac	acc	taa	acc	tta	ccc	14
															Pro	14
	-	35					40			y	****	45	AIG	шеu	FLO	
acc	tac	aat	aac	cac	ctc	tac	aag	caa	atc	tcc	agt	act	tca	aca	aaa	19
															Gly	
	50					55	-				60					
	·															
gcc	agc	aac	gac	aac	cac	tac	ttc	ggc	tac	agc	acc	ccc	tgg	ggg	tat	24
Ala	Ser	Asn	Asp	Asn	His	Tyr	Phe	Gly	Tyr	Ser	Thr	Pro	Trp	Gly	Tyr	
65					70		•			75					80	
					ttc											288
Phe	Asp	Phe	Asn	Arg	Phe	His	Cys	His	Phe	Ser	Pro	Arg	Asp	Trp	Gln	
				85					90					95		
					aat											330
Arg	Leu	Ile	Asn	Asn	Asn	Trp	Gly	Phe	Arg	Pro	Lys	Arg	Leu	Asn	Phe	
			100					105					110			
					caa											384
Lys	Leu		Asn	Ile	Gln	Val		Glu	Val	Thr	Thr	Asn	Asp	Gly	Val	
		115					120					125				
		- - -														
aca mb-	acc	atc	gct	aat	aac	ctt	acc	agc	acg	gtt	caa	gtc	ttc	tcg	gac	432
THE		тте	АТЯ	Asn	Asn		Thr	Ser	Thr	Val		Val	Phe	Ser	Asp	
	130					135					140					

47

_			_		ccg Pro 150		-						_		•	480
		-		_	gcg Ala	-			-		-					528
-	_				ggc Gly	-		-			-					576
-	-	-			cct Pro		•	-		-	-					624
					ttt Phe							_				672
	-	-	-	-	gac Asp 230		-	-					-			720
_			-		aga Arg				-			-	-			768
-	-	_	-		agc Ser	-				•		-		•	-	816
					cct Pro						-	_	_	-		864
					aac Asn										-	912
					aat Asn 310											960
					aaa Lys			-		-				-	-	1008

	att Ile 340							1056
	gtc Val						aac Asn '	1104
	acc Thr							1152
	gac Asp							1200
	gtg Val							1248
	att Ile 420							1296
	ttt Phe							1344
	gtt Val							1392
	ttc Phe					 _	 -	1440
	gag Glu							1488
	aca Thr 500							1536
	aat Asn							1584

tac ctt acc cgt ccc ctg taa Tyr Leu Thr Arg Pro Leu 530 1605

<210> 17 <211> 534

<212> PRT

<213> AAV-1

<400> 17

Met Ala Ser Gly Gly Gly Ala Pro Met Ala Asp Asn Asn Glu Gly Ala 1 5 . 10 . 15

Asp Gly Val Gly Asn Ala Ser Gly Asn Trp His Cys Asp Ser Thr Trp
20 25 30

Leu Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro 35 $4\dot{\nu}$ 45

Thr Tyr Asn Asn His Leu Tyr Lys Gln Ile Ser Ser Ala Ser Thr Gly 50 55 60

Ala Ser Asn Asp Asn His Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr 65 70 75 80

Phe Asp Phe Asn Arg Phe His Cys His Phe Ser Pro Arg Asp Trp Gln 85 90 95

Arg Leu Ile Asn Asn Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe 100 105 110

Lys Leu Phe Asn Ile Gln Val Lys Glu Val Thr Thr Asn Asp Gly Val 115 120 125

Thr Thr Ile Ala Asn Asn Leu Thr Ser Thr Val Gln Val Phe Ser Asp 130 135 140

Ser Glu Tyr Gln Leu Pro Tyr Val Leu Gly Ser Ala His Gln Gly Cys 145 150 155 160

Leu Pro Pro Phe Pro Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr
165 170 175

Leu Thr Leu Asn Asn Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr 180 185 190

Cys Leu Glu Tyr Phe Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe 195 200 205

- Thr Phe Ser Tyr Thr Phe Glu Glu Val Pro Phe His Ser Ser Tyr Ala 210 215 220
- His Ser Gln Ser Leu Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr 225 230 235 240
- Leu Tyr Tyr Leu Asn Arg Thr Gln Asn Gln Ser Gly Ser Ala Gln Asn 245 250 255
 - Lys Asp Leu Leu Phe Ser Arg Gly Ser Pro Ala Gly Met Ser Val Gln 260 265 270
 - Pro Lys Asn Trp Leu Pro Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser 275 280 285
 - Lys Thr Lys Thr Asp Asn Asn Ser Asn Phe Thr Trp Thr Gly Ala 290 295 300
 - Ser Lys Tyr Asn Leu Asn Gly Arg Glu Ser Ile Ile Asn Pro Gly Thr 305 310 315 320
- Ala Met Ala Ser His Lys Asp Asp Glu Asp Lys Phe Phe Pro Met Ser 325 330 335
- Gly Val Met Ile Phe Gly Lys Glu Ser Ala Gly Ala Ser Asn Thr Ala 340 345 350
- Leu Asp Asn Val Met Ile Thr Asp Glu Glu Glu Ile Lys Ala Thr Asn 355 360 365
- Pro Val Ala Thr Glu Arg Phe Gly Thr Val Ala Val Asn Phe Gln Ser 370 380
- Ser Ser Thr Asp Pro Ala Thr Gly Asp Val His Ala Met Gly Ala Leu 385 390 395 400
- Pro Gly Met Val Trp Gln Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile 405 410 415
- Trp Ala Lys Ile Pro His Thr Asp Gly His Phe His Pro Ser Pro Leu 420 425 430
- Met Gly Gly Phe Gly Leu Lys Asn Pro Pro Pro Gln Ile Leu Ile Lys 435 440 445

Asn Thr Pro Val Pro Ala Asn Pro Pro Ala Glu Phe Ser Ala Thr Lys
450 455 460

Phe Ala Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Ser Val Glu 465 470 475 480

Ile Glu Trp Glu Leu Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu
485 490 495

Val Gln Tyr Thr Ser Asn Tyr Ala Lys Ser Ala Asn Val Asp Phe Thr 500 505 510

Val Asp Asn Asn Gly Leu Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg 515 520 525

Tyr Leu Thr Arg Pro Leu 530

<210> 18 <211> 4681 <212> DNA <213> aav-2

<400> 18

tiggccacte cetetege egetegeteg etcactgagg eegggcgace aaaggtegee 60 egacgccegg gettigeeeg ggeggeetea gtgagegage gagegegeag agagggagtg 120 gecaacteca teactagggg treetggagg ggtggagteg tgaegtgaat taegteatag 180 ggttagggag gteetgtatt agaggteaeg tgagtgttt gegacatttt gegacaceat 240 gtggtcacge tgggtattta agecegagtg ageaegeagg gtetecattt tgaageggga 300 ggtttgaacg egeageegee atgeegggt triacgagat tgtgattaag gteeceageg 360 acettgacgg geatetgee ggeatreetg acagetttg gaactgggtg geegagaagg 420 aatgggagt geegeagat tetgacatgg atetgageag geaeeeetga 480 eegtggeega gaagetgeag egegactte tgaeggaatg gegeegtgtg agtaaggeee 540 eggaggeeet triettigg eaattgaga agggagagag etaetteeae atgeaegge 600 tegtggaaac eaceggggtg aaateeatgg tritgggaeg trieetgagt eagattegeg 660 aaaaactgat teagagaatt taeegeggga tegageegae tritgeeaaac tggttegegg 720

tcacaaagac Cagaaatggc gccggaggcg ggaacaaggt ggtggatgag tgctacatcc 780 ccaattactt gctccccaaa acccagcctg agctccagtg ggcgtggact aatatggaac 840 agtatttaag cgcctgtttg aatctcacgg agcgtaaacg gttggtggcg cagcatctga 900 cgcacgtgtc gcagacgcag gagcagaaca aagagaatca gaatcccaat tctgatgcgc 960 cggtgatcag atcaaaaact tcagccaggt acatggagct ggtcgggtgg ctcgtggaca 1020 aggggattac ctcggagaag cagtggatcc aggaggacca ggcctcatac atctccttca 1080 atgcggcctc caactcgcgg tcccaaatca aggctgcctt ggacaatgcg ggaaagatta 1140 tgagcctgac taaaaccgcc cccgactacc tggtgggcca gcagcccgtg gaggacattt 1200 ccagcaatcg gatttataaa attttggaac taaacgggta cgatccccaa tatgcggctt 1260 ccgtctttct gggatgggcc acgaaaaagt tcggcaagag gaacaccatc tggctgtttg 1320 ggcctgcaac taccgggaag accaacatcg cggaggccat agcccacact gtgcccttct 1380 acgggtgcgt aaactggacc aatgagaact ttcccttcaa cgactgtgtc gacaagatgg 1440 tgatctggtg ggaggaggyg aagatgaccg ccaaggtcgt ggagtcggcc aaagccattc 1500 tcggaggaag caaggtgcgc gtggaccaga aatgcaagtc ctcggcccag atagacccga 1560 ctcccgtgat cgtcacctcc aacaccaaca tgtgcgccgt gattgacggg aactcaacga 1620 cettegaaca ceageageeg ttgeaagace ggatgtteaa atttgaacte accegeegte 1680 tggatcatga ctttgggaag gtcaccaagc aggaagtcaa agactttttc cggtgggcaa 1740 aggatcacgt ggttgaggtg gagcatgaat tctacgtcaa aaagggtgga gccaagaaaa 1800 gacccgcccc cagtgacgca gatataagtg agcccaaacg ggtgcgcgag tcagttgcgc 1860 agccatcgac gtcagacgcg gaagcttega tcaactacgc agacaggtac caaaacaaat 1920 gttctcgtca cgtgggcatg aatctgatgc tgtttccctg cagacaatgc gagagaatga 1980 atcagaattc aaatatctgc ttcactcacg gacagaaaga ctgtttagag tgctttcccg 2040 tgtcagaatc tcaaccggtt tctgtcgtca aaaaggcgta tcagaaactg tgctacattc 2100 atcatatcat gggaaaggtg ccagacgctt gcactgcctg cgatctggtc aatgtggatt 2160

tggatgactg catctttgaa caataaatga tttaaatcag gtatggctgc cgatggttat 2220 cttccagatt ggctcgagga cactctctct gaaggaataa gacagtggtg gaagctcaaa 2280 cctggcccac caccaccaaa gcccgcagag cggcataagg acgacagcag gggtcttgtg 2340 cttcctgggt acaagtacct cggacccttc aacggactcg acaagggaga gccggtcaac 2400 gaggcagacg ccgcggccct cgagcacgac aaagcctacg accggcagct cgacagcgga 2460 gacaacccgt acctcaagta caaccacgcc gacgcggagt ttcaggagcg ccttaaagaa 2520 gatacgtctt ttgggggcaa cctcggacga gcagtcttcc aggcgaaaaa gagggttctt 2580 gaacctctcg gcctggttga ggaacctgtt aagacggctc cgggaaaaaa gaggccggta 2640 gagcactete etgtggagee agacteetee tegggaaceg gaaageeggg ceageageet 2700 gcaagaaaaa gattgaattt tggtcagact ggagacgcag actcagtacc tgacccccag 2760 ceteteggae agecaccage agececetet ggtetgggaa etaataegat ggetaeagge 2820 agtggcgcac caatggcaga caataacgag ggcgccgacg gagtgggtaa ttcctccgga 2880 aattggcatt gcgattccac atggatgggc gacagagtca tcaccaccag cacccgaacc 2940 tgggccctgc ccacctacaa caaccacctc tacaaacaaa tttccagcca atcaggagcc 3000 tcgaacgaca atcactactt tygctacagc accccttggg ggtattttga cttcaacaga 3060 ttccactgcc acttttcacc acgtgactgg caaagactca tcaacaacaa ctggggattc 3120 cgacccaaga gactcaactt caacctcttt aacattcaag tcaaagaggt cacgcagaat 3180 gacggtacga cgacgattgc caataacctt accagcacgg ttcaggtgtt tactgactcg 3240 gagtaccage tecegtacgt ecteggeteg gegeateaag gatgeeteec geegtteeca 3300 gcagacgtct tcatggtgcc acagtatgga tacctcaccc tgaacaacgg gagtcaggca 3360 gtaggacgct cttcatttta ctgcctggag tactttcctt ctcagatgct gcgtaccgga 3420 aacaacttta cetteageta eaettttgag gaegtteett teeacageag etaegeteae 3480 agccagagtc tggaccgtct catgaatcct ctcatcgacc agtacctgta ttacttgagc 3540 agaacaaaca ctccaagtgg aaccaccacg cagtcaaggc ttcagttttc tcaggcccca 3600

gccagtgaca ttcgggacca gtctaggaac tggcttcctg gaccctgtta ccgccagcag 3660 cgagtatgaa agacatctgc ggataacaac aacagtgaat actcgtggac tggagctacc 3720 aagtaccacc tcaatggcag agactetetg gtgaateegg ggeeegeeat ggeaageeac 3780 aaggacgatg aagaaaagtt ttttcctcag agcggggttc tcatctttgg gaagcaaggc 3840 tcagagaaaa Caaatgtgaa cattgaaaag gtcatgatta cagacgaaga ggaaatccca 3900 acaaccaatc ccgtggctac ggagcagtat ggttctgtat ctaccaacct ccagagaggc 3960 aacagacaag cagctaccgc agatgtcaac acacaaggcg ttcttccagg catggtctgg 4020 caggacagag atgtgtacct tcaggggccc atctgggcaa agattccaca cacggacgga 4080 cattttcacc cctctccct catgggtgga ttcggactta aacaccctcc tccacagatt 4140 ctcatcaaga acaccccggt acctgcgaat cettcgacca cettcagtgc ggcaaagttt 4200 gcttccttca tcacacagta ctccacggga cacggtcagc gtggagatcg agtgggagct 4260 gcagaacgaa aacagcaaac gctggaatcc cgaaattcag tacacttcca actacaacaa 4320 gtctgttaat cgtggacttt accgtggata ctaatggcgt gtattcagag cctcgcccca 4380 ttggcaccag atacctgact cgtaatctgt aattgcttgt taatcaataa accgtttaat 4440 tcgtttcagt tgaactttgg tctctgcgta tttctttctt atctagtttc catggctacg 4500 tagataagta gcatggcggg ttaatcatta actacaagga acccctagtg atggagttgg 4560 ccactccctc tctgcgcgct cgctcgctca ctgaggccgg gcgaccaaag gtcgcccgac 4620 gcccgggctt tgccccggcg gcctcagtga gcgagcgagc gcgcagagag ggagtgggca 4680 а 4681

<210> 19

<211> 4683

<212> DNA

<213> aav-6

<400> 19

ttggccactc cctctctgcg cgctcgctcg ctcactgagg ccgggcgacc aaaggtcgcc 60

cgacgcccgg gctttgcccg ggcggcctca gtgagcgagc gagcgcgcag agagggagtg 120 gccaactcca tcactagggg ttcctggagg ggtggagtcg tgacgtgaat tacgtcatag 180 ggttagggag gtcctgtatt agaggtcacg tgagtgtttt gcgacatttt gcgacaccat 240 gtggtcacgc tgggtattta agcccgagtg agcacgcagg gtctccattt tgaagcggga 300 ggtttgaacg cgcagcgcca tgccggggtt ttacgagatt gtgattaagg tccccagcga 360 cettgacgag catetgeecg geatttetga cagetttgtg aactgggtgg eegagaagga 420 atgggagttg ccgccagatt ctgacatgga tctgaatctg attgagcagg cacccctgac 480 cgtggccgag aagctgcagc gcgacttcct ggtccactgg cgccgcgtga gtaaggcccc 540 ggaggccctc ttctttgttc agttcgagaa gggcgagtcc tacttccacc tccatattct 600 ggtggagacc acgggggtca aatccatggt gctgggccgc ttcctgagtc agattagcga 660 caagetggtg cagaceatet accgegggat egageegace etgeceaact ggttegeggt 720 gaccaagacg cgtaatggcg ccggaggggg gaacaaggtg gtggacgagt gctacatccc 780 caactacctc ctgcccaaga ctcagcccga gctgcagtgg gcgtggacta acatggagga 840 gtatataagc gcgtgtttaa acctggccga gcgcaaacgg ctcgtggcgc acgacctgac 900 ccacgtcagc cagacccagg agcagaacaa ggagaatctg aaccccaatt ctgacgcgcc 960 tgtcatccgg tcaaaaacct ccgcacgcta catggagctg gtcgggtggc tggtggaccg 1020 gggcatcacc teegagaage agtggateea ggaggaeeag geetegtaea teteetteaa 1080 egeogeetee aastegeggt eccagateaa ggeogetetg gacaatgeeg gcaagateat 1140 ggcgctgacc aaatccgcgc ccgactacct ggtaggcccc gctccgcccg ccgacattaa 1200 aaccaaccgc atttaccgca teetggaget gaacggetac gaccetgeet acgeeggete 1260 cgtctttctc ggctgggccc agaaaaggtt cggaaaacgc aacaccatct ggctgtttgg 1320 gccggccacc acgggcaaga ccaacatcgc ggaagccatc gcccacgccg tgcccttcta 1380 cggctgcgtc aactggacca atgagaactt tcccttcaac gattgcgtcg acaagatggt 1440 gatctggtgg gaggagggca agatgacggc caaggtcgtg gagtccgcca aggccattct 1500

cggcggcagc aaggtgcgcg tggaccaaaa gtgcaagtcg tccgcccaga tcgatcccac 1560 ccccgtgatc gtcacctcca acaccaacat gtgcgccgtg attgacggga acagcaccac 1620 cttcgagcac cagcagccgt tgcaygaccg gatgttcaaa tttgaactca cccqccqtct 1680 ggagcatgac tttggcaagg tgacaaagca ggaagtcaaa gagttettee getgggegca 1740 ggatcacgtg accgaggtgg cgcatgagtt ctacgtcaga aagggtggag ccaacaacag 1800 accegeeece gatgaegegg ataaaagega geeeaagegg geetgeeeet eagtegegga 1860 tccatcgacg tcagacgcgg aaggagctcc ggtggacttt gccgacaggt accaaaacaa 1920 atgttctcgt cacgcgggca tgcttcagat gctgtttccc tgcaaaacat gcgagagaat 1980 gaatcagaat ttcaacattt gcttcacgca cgggaccaga gactgttcag aatgtttccc 2040 cggcgtgtca gaatctcaac cggtcgtcag aaagaggacg tatcggaaac tctgtgccat 2100 teateatetg etggggeggg etecegagat tgettgeteg geetgegate tggteaacgt 2160 ggatctggat gactgtgttt ctgagcaata aatgacttaa accaggtatg gctgccgatg 2220 gttatcttcc agattggctc gaggacaacc tctctgaggg cattcggcag tggtgggact 2280 tgaaacctgg agccccgaaa cccaaagcca accagcaaaa gcaggacgac ggccggggtc 2340 tggtgcttcc tggctacaag tacctcggac ccttcaacgg actcgacaag ggggagcccg 2400 tcaacgcggc ggatgcagcg gccctcgagc acgacaaggc ctacgaccag cagctcaaag 2460 cgggtgacaa tccgtacctg cggtataacc acgccgacgc cgagtttcag gagcgtctgc 2520 aagaagatac gtcttttggg gycaacctcg ggcgagcagt cttccaggcc aagaagaggg 2580 ttctcgaacc ttttggtctg gttgaggaag gtgctaagac ggctcctgga aagaaacgtc 2640 cggtagagca gtcgccacaa gagccagact cctcctcggg cattggcaag acaggccagc 2700 agcccgctaa aaagagactc aattttggtc agactggcga ctcagagtca gtccccgacc 2760 cacaacctct cggagaacct ccagcaaccc ccgctgctgt gggacctact acaatqqctt 2820 caggcggtgg cgcaccaatg gcagacaata acgaaggcgc cgacggagtg ggtaatgcct 2880 caggaaattg gcattgcgat tccacatggc tgggcgacag agtcatcacc accagcaccc 2940

gaacatgggc cttgcccacc tataacaacc acctctacaa gcaaatctcc agtgcttcaa 3000 cgggggccag caacgacaac cactacttcg gctacagcac cccctggggg tattttgatt 3060 tcaacagatt ccactgccat ttctcaccac gtgactggca gcgactcatc aacaacaatt 3120 ggggattccg gcccaagaga ctcaacttca agctcttcaa catccaagtc aaggaggtca 3180 cgacgaatga tggcgtcacg accatcgcta ataaccttac cagcacggtt caagtcttgt 3240 cggactcgga gtaccagttc ccgtacgtcc tcggctctgc gcaccagggc tgcctccctc 3300 cgttcccggc ggacgtgttc atgattccgc agtacggcta cctaacgctc aacaatggca 3360 gccaggcagt gggacgctca tccttttact gcctggaata tttcccatcg cagatgctga 3420 gaacgggcaa taactttacc ttcagctaca ccttcgagga cgtgcctttc cacagcagct 3480 acgcgcacag ccagagcctg gaccggctga tgaatcctct catcgaccag tacctgtatt 3540 acctgaacag aactcacaat cagtccggaa gtgcccaaaa caaggacttg ctgtttagcc 3600 gtgggtctcc agctggcatg tctgttcagc ccaaaaactg gctacctgga ccctgttacc 3660 ggcagcagcg cgtttctaaa acaaaaacag acaacaacaa cagcaacttt acctggactg 3720 gtgcttcaaa atataacctt aatgggcgtg aatctataat caaccctggc actgctatgg 3780 cctcacacaa agacgacaaa gacaagttet tteecatgag eggtgteatg atttttggaa 3840 aggagagege eggagettea aacaetgeat tggacaatgt catgateaca gacgaagagg 3900 aaatcaaage cactaaccce gtggccaccy aaagatttgg gactgtggca gtcaatctcc 3960 agagcagcag cacagaccct gcgaccggag atgtgcatgt tatgggagcc ttacctggaa 4020 tggtgtggca agacagagac gtatacctgc agggtcctat ttgggccaaa attcctcaca 4080 cggatggaca ctttcacccg tetectetca tgggcggett tggaettaag cacccgcete 4140 ctcagatect catcaaaaac acgeetgtte etgegaatee teeggeagag tttteggeta 4200 caaagtttgc ttcattcatc acccagtatt ccacaggaca agtgagcgtg gagattgaat 4260 gggagctgca gaaagaaaac aycaaacgct ggaatcccga agtgcagtat acatctaact 4320 atgcaaaatc tgccaacgtt gatttcactg tggacaacaa tggactttat actgagcctc 4380

<210> 20

<211> 16

<212> DNA

<213> rep binding motif

<400> 20

gctcgctcgc tcgctg

16

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT) (51) International Patent Classification 7: (11) International Publication Number: WO 00/28061 **A3** C12N 15/86, 15/35, 5/10, A61K 48/00 (43) International Publication Date: 18 May 2000 (18.05.00) (21) International Application Number: PCT/US99/25694 (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, (22) International Filing Date: GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, 2 November 1999 (02.11.99) KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, (30) Priority Data: SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, 60/107,114 5 November 1998 (05.11.98) US ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, (71) Applicant (for all designated States except US): THE DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), TRUSTEES OF THE UNIVERSITY OF PENNSYLVA-OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, NIA [US/US]; Suite 300, 3700 Market Street, Philadelphia, MR, NE, SN, TD, TG). PA 19104-3147 (US). Published (72) Inventors; and (75) Inventors/Applicants (for US only): WILSON, James, M. With international search report. [US/US]; 1350 N. Avignon Drive, Gladwyne, PA 19035 (US). XIAO, Weidong [CN/US]; Apartment P4, 155 Wash-(88) Date of publication of the international search report: ington Lane, Jenkintown, PA 19046 (US). 3 August 2000 (03.08.00) (74) Agents: KODROFF, Cathy, A. et al.; Howson & Howson, Spring House Corporate Center, P.O. Box 457, Spring House, PA 19477 (US).

(54) Title: ADENO-ASSOCIATED VIRUS SEROTYPE I NUCLEIC ACID SEQUENCES, VECTORS AND HOST CELLS CONTAINING SAME

(57) Abstract

The nucleic acid sequences of adeno-associated virus (AAV) serotype 1 are provided, as are vectors and host cells containing these sequences and functional fragments thereof. Also provided are methods of delivering genes via AAV-1 derived vectors.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
АT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Мопасо	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	T.J	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	•
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Trinidad and Tobago
BR	Brazil	IL	Israel	MR	Mauritania	UG	Ukraine
BY	Belarus	IS	Iceland	MW	Malawi	US	Uganda
CA	Canada	IT	Italy	MX	Mexico		United States of America
CF	Central African Republic	JP	Japan	NE	Niger	UZ VN	Uzbekistan
CG	Congo	KE	Kenya	NL	Netherlands		Viet Nam
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	YU	Yugoslavia
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand	ZW	Zimbabwe
СМ	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	- · -		
CU	Cuba	KZ	Kazakstan	RO	Portugal Romania		•
CZ	Czech Republic	LC	Saint Lucia	RU			
DE	Germany	LÍ	Liechtenstein		Russian Federation		
DK	Denmark	LK	Sri Lanka	SD	Sudan		
EE	Estonia	LR	Liberia	SE	Sweden		
		LR	LIDENA	SG	Singapore		

'INTERNATIONAL SEARCH REPORT

Inti. Jonal Application No PCT/US 99/25694

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/86 C12N15/35 C12N5/10 A61K48/00 According to International Patent Classification (IPC) or to both national classification and IPC **B. FIELDS SEARCHED** Minimum documentation searched (classification system followed by classification symbols) IPC 7 C12N A61K Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used) C. DOCUMENTS CONSIDERED TO BE RELEVANT Category Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No. RUTLEDGE E. A. ET AL.: "Infectious clones 1-23 and vectors derived from adeno-associated virus (AAV) serotypes other than AAV type 2." JOURNAL OF VIROLOGY, vol. 72, no. 1, January 1998 (1998-01), pages 309-319, XP002137089 ISSN: 0022-538X cited in the application the whole document WO 98 11244 A (SAFER BRIAN ;US HEALTH 1-23 (US); CHIORINI JOHN A (US); KOTIN ROBERT M) 19 March 1998 (1998-03-19) the whole document X Further documents are listed in the continuation of box C. Patent family members are listed in annex. Special categories of cited documents : "T" later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the International "X" document of particular relevance; the claimed invention filing date cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone 1. document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such docu-*O* document referring to an oral disclosure, use, exhibition or ments, such combination being obvious to a person skilled in the art. document published prior to the international filing date but later than the priority date claimed "&" document member of the same patent family Date of the actual completion of the international search Date of mailing of the international search report 8 May 2000 22/05/2000 Name and mailing address of the ISA **Authorized officer** European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016 Mandl, B



Int Bonal Application No PCT/US 99/25694

C (C	Man) POCINENTS CONSIDERATE TO THE	PCT/US 99/25694					
Category •	ation) DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages		In-leading to the N				
	от по терезадов	_ ·	Relevant to claim No.				
°,X	XIAO W. ET AL.: "Gene therapy vectors based on adeno-associated virus type 1." JOURNAL OF VIROLOGY, vol. 73, no. 5, May 1999 (1999-05), pages 3994-4003, XP002137090 ISSN: 0022-538X the whole document		1-23				
			·				



INTERNATIONAL SEARCH REPORT

...emational application No.

PCT/US 99/25694

Box I	Observations where certain claims were found unsearchable (Continuation of Item 1 of Ifret sheet)
This Inte	emational Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
1. X	Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely: Remark: Although claims 18-20 and 22, as far as an in vivo application is concerned, are directed to a method of treatment of the human or animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2.	Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3.	Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II	Observations where unity of invention is lacking (Continuation of item 2 of first sheet)
This inte	emational Searching Authority found multiple inventions in this international application, as follows:
1.	As all required additional search fees were timely paid by the applicant, this international Search Report covers all searchable claims.
2.	As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3.	As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4.	No required additional search fees were timely paid by the applicant. Consequently, this international Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
Remark	on Protest The additional search fees were accompanied by the applicant's protest. No protest accompanied the payment of additional search fees.





int	donal	Application No
PCT.	/US	99/25694

Information on patent family members					Int Bonal Application No PCT/US 99/25694			
Patent document cited in search report		Publication date	Patent family member(s)			Publication date		
WO 9811244	A	19-03-1998	AU EP	46456 09326	97 A 94 A	02-04-1998 04-08-1999		
	•							